

lwi-Māori Partnership Board Health Profile:

Te Moana a Toi



Iwi-Māori Partnership Board Health Profile: Te Moana a Toi

Volume One

Recommended citation: Curtis, E., Loring, B., Walker, R., Pearse, T., Gilbert-Perenise, S., Gray, G., Akuhata-Huntington, Z., Latham, K., Kiriona, K. (2023). *Iwi-Māori Partnership Board Health Profile: Te Moana a Toi. Volume One.* Te Aka Whai Ora – Māori Health Authority; Auckland.

15th December 2023

ISBN - 978-1-991286-03-1



Ngā kupu whakamihi Acknowledgements

Toitū te mauri nui

Toitū te mauri roa

Toitū te mauri ora

Tīhei Te Aka Whai Ora e!

Kei ngā whakatiketike ki te rangi, kei ngā whakatamarahi ki te whenua, koutou e kōkiri ana i te Pae Ora nō roto mai i te oranga nui, te oranga roa o ō tātou whānau, hapū, iwi puta i Aotearoa whānui – tēnā koutou!

E pēnei ana te nui, me te hari o ngā mihi ki a koutou e ngā kaiwhakairo i te tatauranga Hauora Māori kia pai ai te whakatakoto kupu mō tā tātou kaupapa, mō Te Aka Whai Ora.

Kāore e ārikarika nei ngā mihi nui ki a koutou -

huri noa, tēnā koutou, tēnā koutou, tēnā tātou katoa.

Ngā kupu whakamihi - Acknowledgements

Many people have contributed their time and expertise to the lwi-Māori Partnership Board health profiles.

These profiles have been written by Elana Curtis, Belinda Loring, Rhiannon Walker, Tyla Pearse, Sade Gilbert-Perenise, George Gray, with overall leadership and direction provided by Zaine Akuhata-Huntington, Kadin Latham and Kingi Kiriona. This work has been supported by a wider team at Te Aka Whai Ora, including Nigel Chee, Paula Searle, Teei Kaiaruna, Riki Kyle, Yahp Jasperse, Swathy Babu Kathanaparambil, Ngarangi Williams, Michele Bristow, Ngaringi Katipa, TeMatetahuna Paki, and Raniera Albert.

We thank Doone Winnard, Delwyn Armstrong and Gary Jackson at Te Whatu Ora for their leadership and support with data quality assurance.

We are also very grateful to the following people who assisted us to obtain and analyse data: Sydney Kingstone, Mildred (Ai Wei) Lee, Michael Walsh, Dean Papaconstantinou; Liam Gray from Statistics New Zealand; and Kirk Paterson from Manatū Hauora.

Shelley Cousins from RUN developed the document template and graphic design.

We would like to thank Suzanne Pitama and Reigna Morgan for providing helpful peer review of earlier drafts.

We acknowledge the leadership of Bridget Robson and colleagues at Te Rōpū Rangahau Hauora a Eru Pōmare, University of Otago Wellington, whose earlier work on the 2015 District Health Board Māori Health Profiles was used as a basis for these Iwi-Māori Partnership Board profiles.



Te kupu takamua Foreword

Te kupu takamua - Foreword

We are extremely pleased to present this report that provides the most up-to-date snapshot of Māori health for the newly formed lwi-Māori Partnership Boards.

In doing so, we acknowledge the legacy of work associated with Māori-led health data reporting to date – from the seminal *Hauora* series to *Tatau Kahukura* and the *2015 District Health Board Māori Health Profiles*, this report continues the commitment to excellence that Māori communities and whānau both need and deserve.

Iwi-Māori Partnership Boards were created under the Pae Ora (Healthy Futures) Act 2022 to provide a vehicle for local feedback and leadership on how the health sector is performing to meet the needs and aspirations of whānau in their area. Iwi-Māori Partnership Boards have a pivotal role to play in determining how health services and public health interventions should be designed and delivered.

Te Aka Whai Ora welcomes the contribution of each Iwi-Māori Partnership Board to use the data presented in these reports to understand what issues are important to them and what response(s) are needed to ensure their tino rangatiratanga and mana motuhake over their health and wellbeing are being realised. The data presented in this profile require contextualisation - they are a starting point for Iwi-Māori Partnership Boards to interpret, together with other sources of information, and decide how best to respond to the needs (and rights) of the whānau within their rohe.

This report represents the first wave of analysis (Volume One). This volume includes key demographic information, mauri ora (overall health status), whānau ora (healthy families) and wai ora (healthy environments) indicators specific to each Iwi-Māori Partnership Board. A second volume with additional indicators focused on Te Aka Whai Ora-identified health priority areas (e.g. cancer, long-term conditions, first 1,000 days and mental health) will be released early in 2024.

The data presented within these profiles are a dimension of 'whānau voice'. They represent Māori stories and Māori lived experience and should be valued as a taonga for the health system to use and respond to as part of the broader commitment to Te Tiriti o Waitangi and equity.

We are extremely humbled by the sacrifices that have been made by our people: externally, as Iwi-Māori Partnership Boards have been established, and within the organisation, to produce this output in such a short time-frame since our establishment as an entity in July 2022.

We thank our partners who have contributed to this report and hope that this commitment to excellence in Māori health continues - mō āke tonu atu.

Ngā mihi,

Tipa Mahuta

Waikato, Maniapoto, Ngāpuhi

Te Kaihautū (Chair)

Pu (

Riana Manuel

Ngāti Pukenga, Ngāti Maru, Ngāti Kahungunu

Te Aka Matua (Chief Executive)





Te ihirangi - Table of Contents

Ngā ku	upu whakamihi - Acknowledgements	4
Te kup	ou takamua - Foreword	6
Te ihira	angi - Table of Contents	8
List of	Tables	10
List of	Figures	12
List of	Abbreviations, Acronyms and Initialisms	13
Māori (Glossary	14
1. Te	e kupu whakataki - Introduction	16
1.1.	Overview of Iwi-Māori Partnership Boards	16
1.2.	Purpose and audience for this report	18
1.3.	Positioning	18
1.4.	Understanding Māori health and health inequities	18
1.5.	Scope for these profiles	20
1.6.	Data sources	22
1.7.	How to understand this report	22
2. No	gā tatauranga taupori matua - Key demographics	24
2.1.	About Te Moana a Toi	24
3. M	auri ora - overall health status	28
3.1.	Life Expectancy	28
3.2.	Self-assessed health	29
3.3.	Mortality	30
4. W	/hānau ora - Healthy families	36
5. W	ai ora - Healthy environments	40
5.1.	Education	40
5.2.	Work	40
5.3.	Income and standard of living	43
5.4.	Housing	45
5.5.	Primary Care Enrolment	47
Appen	dix 1: IMPB Māori population projections	49
Appen	dix 2: Technical notes	56
1. E	xplanation of statistical terms used in this report	56
95	5% confidence interval	56

Age standardisation	57
Rate ratios	58
Rate difference	59
2. Key methods and quality limitations of key data sources	60
Numerators	60
Denominators	60
Ethnicity data	60
Age-standardised and crude rates	61
Confidence intervals	61
Rate ratios	61
Demography data	62
The Census of Population and Dwellings	62
Geographical alignment between IMPB and DHB areas	64
Life expectancy	65
Mortality data	65
Te Kupenga Survey	65
Primary care enrolment	66
NZ Index of Deprivation 2018	66
Geographic Classification of Health	66
Appendix 3: ICD-10-AM Codes	68
Appendix 4: Māori 2001 Population	70
Te rārangi tohutoro - References	72



List of Tables

Table 1 - Population estimates by age group, Te Moana a Toi, 202325
Table 2 - Population projections, Te Moana a Toi, 2023 to 204325
Table 3 - Life expectancy at birth, Māori and non-Māori, Te Moana a Toi, 2018 to 202228
Table 4 - Decomposition of the ethnic gap in life expectancy by avoidable category - Māori compared with non-Māori/non-Pacific, 2018 to 2020, Te Manawa Taki Region29
Table 5 - Health status reported by Māori aged 15 years and over, Te Moana a Toi, 201829
Table 6 - Leading causes of death for Māori, all ages, Bay of Plenty DHB, 2014 to 201830
Table 7 - Leading causes of death for Māori, all ages, Aotearoa, 2014 to 201831
Table 8 - All-cause deaths, all ages, Bay of Plenty DHB, 2014 to 2018
Table 9 - Potentially avoidable deaths, ages 0-74 years, Bay of Plenty DHB, 2014 to 201832
Table 10 - Leading causes of potentially avoidable mortality, ages 0-74 years, Bay of Plenty DHB 2014 to 2018
Table 11 - Leading causes of potentially avoidable mortality, ages 0-74 years, Aotearoa, 2014 to 2018
Table 12 - Whānau well-being reported by Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018
Table 13 - Whānau composition reported by Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018
Table 14 - Access to whānau support, Māori aged 15 years and over, Te Moana a Toi and Aotearoa 2018
Table 15 - Importance of Māori culture and spirituality, Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018
Table 16 - Use of te reo Māori in the home, Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018
Table 17 - Access to marae, Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018 38
Table 18 - Māori aged 15 years and over who took part in traditional healing or massage in last 12 months, Te Moana a Toi and Aotearoa, 2018
Table 19 - Adults aged 20 years and over with a Level 2 Certificate or higher, Bay of Plenty DHB 201840
Table 20 - Labour force status, 15 years and over, Bay of Plenty DHB, 201840
Table 21 - Leading industries in which Māori were employed, Bay of Plenty DHB, 201841
Table 22 - Leading occupations in which Māori were employed, Bay of Plenty DHB, 201842
Table 23 - Unpaid work, 15 years and over, Bay of Plenty DHB, 201843
Table 24 - Unmet needs reported by Māori aged 15 years and over to keep costs down in the last 12 months. Te Moana a Toi and Aotearoa. 2018



able 25 - People 20 years and over whose total annual personal income in \$20,000 or less, Ba lenty DHB, 2018	•
able 26 - People with no access to a motor vehicle, Bay of Plenty DHB, 2018	44
able 27 - People with no access to telecommunications, Bay of Plenty DHB, 2018	45
able 28 - Housing tenure, 20 years and over, Bay of Plenty DHB, 2018	45
able 29 - People living in crowded households (requiring at least one more bedroom), Bay of Pl HB, 2018	•
able 30 - People experiencing housing quality issues sometimes or always, Bay of Plenty D	
able 31 - People living in households where there is no source of heating, Bay of Plenty DHB, 2	
able 32 - People enrolled with primary care, Bay of Plenty DHB, October 2023	
able 33 - Māori population projections, single year, Bay of Plenty DHB, by 5-year age band, 201	
able 34 - Quality ratings 2018 Census variables included in this report	63
able 35 - Potentially avoidable mortality ICD-10-AM codes	68
able 36 - 2001 Census total Māori population	70



List of Figures

Figure 1 - Map of Iwi-Māori Partnership Board areas	17
Figure 2 - Te Kupenga Hauora Māori modified model for explaining Indigenous/ethnic determinar of health	
Figure 3 - Map of Te Moana a Toi IMPB with DHB boundaries, 2023	24
Figure 4 - Population distribution by urban and rural classification, Bay of Plenty DHB, 2023	26
Figure 5 - NZDep2018 distribution of Māori and non-Māori by decile. Bay of Plenty DHB 2018	43



List of Abbreviations, Acronyms and Initialisms

ANZSCO	Australian and New Zealand Standard Classification of Occupations
ANZSIC	Australian and New Zealand Standard Industrial Classification
Av	Average
CI	Confidence Intervals
COPD	Chronic Obstructive Pulmonary Disease
DHB	District Health Board
ERP	Estimated resident population
GCH	Geographic Classification for Health
ICD-10-AM	International Statistical Classification of Diseases and Related Health Problems, Tenth Revision, Australian Modification
IMPB	lwi-Māori Partnership Board
NHI	National Health Index
No	Number
NZ	Aotearoa/New Zealand
NZDep2018	New Zealand Index of Deprivation 2018
PHO	Primary Health Organisation
RR	Rate ratio
SA1	Statistical Area Level 1
SA2	Statistical Area Level 2
StatsNZ	Statistics New Zealand
TKHM	Te Kupenga Hauora Māori
UR	Usually resident
WHO	World Health Organization



Māori Glossary

Aotearoa	New Zealand
Hāpori Māori	Māori communities
Hauora Māori	Māori health
Hui	Meeting, gathering
lwi	Tribe
Kaupapa Māori	Māori initiative, approach, topic, agenda, principle, ideology
Manatū Hauora	Ministry of Health
Māori	Indigenous people(s) of Aotearoa New Zealand
Marae	Complex of buildings significant to Māori, may include, but not limited to, wharenui, wharekai, and urupā
Mauri ora	Overall health status
Mō āke tonu atu	Forever
Ngā āpitihanga	Appendices
Ngā kupu whakamihi	Acknowledgements
Ngā mihi	Greetings
Ngā tatauranga taupori matua	Key demographics
Pae ora	Healthy futures
Rohe	Region
Tangi	Funeral, mourning
Taonga	Treasure
Tatau Kahukura	Māori Health Chartbook 2015
Te Aka Whai Ora	Māori Health Authority
Te ihirangi	Contents
Te Kupenga Hauora Māori	Department of Māori Health, Faculty of Medical and Health Sciences, The University of Auckland
Te kupu takamua	Foreword
Te kupu whakataki	Introduction
Te rārangi tohutoro	References
Te Rōpū Rangahau Hauora a Eru Pōmare	Eru Pomare Māori Health Research Centre, The University of Otago
Te Tiriti o Waitangi	Treaty of Waitangi
Te Whatu Ora	Health New Zealand
Wai ora	Healthy environments
Whakamaua	Māori Health Action Plan: 2020-2025
Whānau	
	Family



1. Te kupu whakataki - Introduction

1.1. Overview of Iwi-Māori Partnership Boards

One of the three purposes of the Pae Ora (Healthy Futures) Act 2022 (Pae Ora) is to "achieve equity in health outcomes among New Zealand's population groups, including by striving to eliminate health disparities, in particular for Māori". Iwi-Māori Partnership Boards (IMPBs) are an important legislated mechanism for the Crown to give effect to the principles of Te Tiriti o Waitangi (the Treaty of Waitangi). The Pae Ora Act requires Health New Zealand (Te Whatu Ora) and the Māori Health Authority (Te Aka Whai Ora) to engage with IMPBs.

The purpose of IMPBs is to represent local Māori perspectives on:

- a) the needs and aspirations of Māori in relation to hauora Māori outcomes; and
- b) how the health sector is performing in relation to those needs and aspirations; and
- c) the design and delivery of services and public health interventions within localities.

The Pae Ora Act sets out the criteria for recognition of an organisation as an IMPB. The criteria ensure the Boards are broadly representative of all Māori within the relevant area and include;

- a) that the proposed boundaries of the area covered by the organisation do not overlap with the boundaries of any area covered by any other IMPB;
- b) that the organisation has taken reasonable steps to engage with relevant Māori communities and groups; and
- the organisation must demonstrate that it has the capacity and capability to perform the necessary functions of IMPBs as set out in the Act, and that the organisation can represent and be accountable to hāpori Māori (Māori communities).

Once the Board of Te Aka Whai Ora is satisfied that an organisation has met the criteria for recognition, they advise the Minister of Health who then recommends the making of an Order in Council so that the organisation can be listed as an IMPB (under Schedule 4 of the Pae Ora Act). On the advice of the Te Aka Whai Ora Board, the Minister of Health can also recommend an Order in Council to vary or remove an IMPB from Schedule 4 of the Pae Ora Act. An important feature of IMPBs is that they can renegotiate boundaries between each other as and when works for the collective. Such is the case for any emerging organisation who must consult with neighbouring IMPBs should their intended boundary result in overlap. This ensures the self-determination of communities, and strategic alignment with community need.

As at July 2023, 15 IMPBs were listed in Schedule 4, as shown in Figure 1.



Figure 1 - Map of Iwi-Māori Partnership Board areas



1.2. Purpose and audience for this report

Under the Pae Ora Act, Te Aka Whai Ora must take reasonable steps to support IMPBs to achieve their purpose, including by providing administrative, analytical, or financial support where needed; and providing sufficient and timely information. These data profiles have been prepared for each IMPB formed in 2023, as part of a commitment by Te Aka Whai Ora to provide IMPBs with health information to inform priorities and actions.

Te Aka Whai Ora has produced these profiles, together with support from Te Whatu Ora, to provide IMPBs with a baseline snapshot of the health of Māori in their rohe (region). These profiles are limited to the data sources and indicators currently available in the government health system, and may not capture all aspects of hauora Māori, determinants of wellbeing, or government responsibility.

1.3. Positioning

This profile has been drafted from a Kaupapa Māori research and epidemiology positioning (Simmonds, Robson et al. 2008). This positioning includes:

- a commitment to high quality ethnicity data reporting and analysis (that includes understanding how ethnicity data are collected and recorded and the implications of these factors on data quality from various sources);
- a commitment to using appropriate comparator groupings (or not) within ethnic data comparisons (that reflect Te Tiriti o Waitangi/rights-based and equity appropriate interpretations) (Harris, Paine et al. 2022), and;
- a strengths-based interpretation of data that rejects 'victim-blame' or 'cultural-deficit' interpretations of any data presented (Curtis 2016).

It is important to note that the identification of inequities between Māori and non-Māori is not a signal of Māori failure or shortcomings. Rather, a Kaupapa Māori positioning foregrounds racism, privilege and power imbalances as the fundamental drivers of ethnic inequities in health for Māori compared to non-Māori (Curtis, Jones et al. 2023).

The data presented in this profile require contextualisation - they are a starting point for IMPBs to interpret, together with other sources of information, and decide how best to respond to the needs (and rights) of their specific population. Although quantitative in nature, the data presented within these profiles are a dimension of 'whānau voice'. They represent Māori stories and Māori lived experience and should be valued as a taonga for the health system to use and respond to as part of the broader commitment to Te Tiriti o Waitangi and equity.

1.4. Understanding Māori health and health inequities

It is important to have a common understanding on what the fundamental drivers or Māori health and health inequities are in order to respond appropriately. A helpful framework is the 'Te Kupenga Hauora Māori (TKHM) modified model' (Curtis, Jones et al. 2023) - a Māori model that draws upon international theorisation on the causation of ethnic health inequities (Figure 2). The TKHM modified model outlines a framework to understand the causes of Māori:non-Māori health inequities within an Aotearoa and Indigenous specific context.


The framework emphasises the importance of distinguishing basic causes from surface (or intervening causes). Overall, changes in basic causes create important changes in health outcomes. Social status categories are created, and reinforced, by basic causes. Social status categories considered to have particular relevance to Māori health outcomes include: ethnicity, socio-economic status, gender, age, and poor health status. In the TKHM modified model, surface causes represent

a number of intervening mechanisms that link *social status* categories such as *ethnicity*, to *health outcomes*. Important intervening mechanisms include: *stress*, *socio-economic opportunities*, *societal resources*, *health services* and *social transmission*. Health *outcomes* reflect the mechanisms by which differences in health status and therefore health inequities are observed or measured. For example, health can vary with respect to *morbidity* (ill health), *mortality* (death rates), presence or absence of *disability*, *mental health* and generalised *wellbeing*.

The TKHM modified model foregrounds colonisation as a key determinant of health inequities underpinning all levels from *basic* to *surface* causes. In doing so, the model acknowledges the historical trauma of colonisation whilst also foregrounding the ongoing contemporary effects of colonisation in today's society. It is not a simple, unidirectional relationship between causes at different levels - but rather there is a dynamic interplay between causes and pathways. Worldviews and positioning are also a basic cause, and privilege alongside racism plays a causative role in Māori health inequities.

Explanations define solutions. Therefore, a conceptual framework can support the understanding of fundamental causes of Indigenous and Māori health inequities and how best to respond to those inequities once they have been identified. Many of the routine data that are collected and reported in Aotearoa, including in this report, focus on the downstream surface causes. It is important to understand that many of these indicators are outcomes/consequences of structural processes of marginalisation that we do not properly measure, and that intervention needs to occur upstream to achieve health equity for Māori.

Figure 2 - Te Kupenga Hauora Māori modified model for explaining Indigenous/ethnic determinants of health



Source: Curtis, Jones et al., 2023

1.5. Scope for these profiles

These profiles are the first reports which specifically focus on data related to IMPBs. These profiles focus on key population demographic data, indicators reflecting key socio-economic determinants of wellbeing, health status and health services indicators. Not every health issue or determinant is included. These IMPB profiles are presented in two volumes:

- Volume One contains key demographic data and projections, overall life expectancy and health outcomes measures, and indicators relating to whānau wellbeing and socio-economic and environmental determinants of wellbeing.
- Volume Two contains health service utilisation and outcomes measures, with a focus on the four health priority areas identified in the 2022 Te Aka Whai Ora Māori Health Priorities Report (Curtis E, Loring B et al. 2022): the first 1000 days, cancer, long term conditions, and mental health and addiction.

These reports are by no means exhaustive, and IMPBs may wish to also refer to other sources of information available through respective government agencies for more in-depth data related to areas such as education, social development, environment, employment or housing. We are limited to currently available data, which may not reflect all indicators of importance to IMPBs, and not all data (for example, on uncommon health conditions) can be meaningfully disaggregated by ethnicity to the level of IMPBs. These IMPB profiles are intended to be used in conjunction with other sources of publicly available health system reporting by the Ministry of Health, Te Whatu Ora, the Health Quality and Safety Commission, Statistics New Zealand (StatsNZ) and other agencies.

There have also been a number of previous sources of reporting specifically on Māori health, which IMPBs may wish to refer to for additional information relevant to their area, including trends over time. Some of these key sources include:

Whakamaua Dashboard¹

This online dashboard presents quantitative measures which assess system performance against the four objectives of Whakamaua: Māori Health Action Plan 2020-2025. From 2023, the Whakamaua dashboard contains some indicators disaggregated by Iwi-Māori Partnership Boards (IMPB). These data for IMPBs use the Health Service Utilisation population as the denominator, which differs slightly from the Census population denominator chosen in these IMPB profiles. The Whakamaua dashboard compares Māori data to non-Māori non-Pacific data.

WAI 2575 Māori Health Trends Report²

This report was compiled by the Ministry of Health in 2019, to inform the Wai 2575 Health Services and Outcomes Kaupapa Inquiry (Wai 2575). The report shows changes of Māori health over the years 1990-2015. Most data are presented at a national level, for Māori compared to non-Māori, and Māori compared to non-Māori non-Pacific, although some variables are available at a District Health Board (DHB) level.

² https://www.health.govt.nz/publication/wai-2575-maori-health-trends-report



¹ https://minhealthnz.shinyapps.io/WhakamauaDashboard/

A Window on the Quality of Aotearoa New Zealand's Health Care 2019 - a view on Māori health equity³

A Window on the Quality of Aotearoa New Zealand's Health Care 2019 - a view on Māori health equity was compiled by the Health Quality and Safety Commission and highlights a number of areas where change is needed in the health system. The report is divided into three chapters. The first analyses inequity between how Māori and non-Māori access and receive health services, and the effects on equity of improvement activities in our system. The second chapter asks why these inequities exist, and the third chapter addresses opportunities for improvement.

• 2015 District Health Board Māori Health Profiles⁴

The 2015 District Health Board Māori Health Profiles were produced by Te Rōpū Rangahau Hauora a Eru Pōmare at the University of Otago in Wellington. The District Health Board Māori Health Profiles present a snapshot of Māori health compared with non-Māori across a range of health and disability-related indicators. They can create a picture of the health status of a DHB's population at a given time and allow some comparison of trends over time. The profiles are available as word and pdf documents, and Excel tables containing data from the profiles together with national rates for most indicators.

• Tatau Kahukura: Māori health statistics⁵

Statistical profiles on Māori health compiled by the Ministry of Health, most recently completed in 2015. Presents Māori compared to non-Māori national level data for a range of health indicators (socio-economic determinants, risk factors, health services and health outcomes), and data are age-standardised to the 2001 Māori population.

Hauora: Māori Standards of Health IV: A study of the years 2000-2005⁶

Hauora: Māori Standards of Health IV, published in 2007, is the most recent edition in the Hauora series, produced by Te Rōpū Rangahau Hauora a Eru Pōmare, and covers the period 2000 to 2005. Careful consideration has been given to the manner in which evidence has been presented and the commentaries are rightly written from Māori perspectives. The first three chapters situate health statistics within the broader context, including the theoretical, demographic and socio-economic contexts. This is followed by chapters on mortality, public hospitalisations, cancer and mental health. This volume of Hauora also includes a number of topic-based chapters from invited authors, including chapters on cardiovascular disease; diabetes; respiratory disease; oral health; disability; sleep problems; occupational safety and health; health in prisons; and the National Primary Medical Care Survey.

To maximise consistency and make it easier for IMPBs to assess how various indicators in their rohe are tracking over time, we have endeavoured to replicate the scope and approach taken in the 2015 District Health Board Māori Health profiles as closely as possible. There are some minor variations in statistical methods, definitions and geographical boundaries for some indicators, which mean that exact comparison with these earlier profiles is not always possible.



³https://www.hqsc.govt.nz/resources/resource-library/a-window-on-the-quality-of-aotearoa-new-zealands-health-care-2019-a-view-on-maori-health-equity-2/

⁴https://www.health.govt.nz/publication/dhb-maori-health-profiles

⁵https://www.health.govt.nz/our-work/populations/maori-health/tatau-kahukura-maori-health-statistics

⁶https://www.otago.ac.nz/wellington/departments/publichealth/research-groups-in-the-department-of-publichealth/erupomare/research/hauora-maori-standards-of-health-iv-a-study-of-the-years-2000-2005

1.6. Data sources

The data presented in this report come from routinely collected national government health datasets and routine national surveys. The main data sources for this report are:


- The 2018 Census of Population and Dwellings
- Te Kupenga 2018 (the Māori Social Survey)
- Mortality registrations
- Te Whatu Ora Primary Care Enrolment data

Data are presented for Māori and non-Māori residents, using the geographical boundaries in each dataset which most closely correspond to the boundaries of the IMPB. For some measures, the closest available match at this time has been the boundaries of the former DHBs covering the IMPB rohe. Where an IMPB area encompasses more than one former DHB, data are presented separately for each DHB area, to provide a sense of variation for Māori within the IMPB.

1.7. How to understand this report

The technical appendix at the end of this report contains further information to help users interpret the data presented. This includes a basic explanation of how to interpret the graphs and tables provided. There is also a description of key methods, including age-standardisation, comparator groups and statistical calculations. The appendix also contains a description of the quality of ethnicity data in each data source used in this profile, and how this may affect the accuracy of information for Māori. Further technical details are provided about the methods and data sources used to compile these reports, so that the methods can be replicated by others.





2. Ngā tatauranga taupori matua - Key demographics

2.1. About Te Moana a Toi

Te Moana a Toi IMPB is home to an estimate of 68,000 Māori in 2023 (Table 1) and consists of parts of the geographic area of the former Bay of Plenty DHB. As Figure 3 shows, not all of the former Bay of Plenty DHB is included in the IMPB health planning area. This report includes data which is specific to the IMPB health planning area where possible, but for indicators where this level of data is not available, data for the entire Bay of Plenty DHB is provided. In this report, where data is presented for the IMPB, it has been mapped to SA2 geographic areas, and where data has been presented for the DHB, it is mapped to DHB boundaries. See the Technical Appendix at the end of this report for more details about how the geographic areas for the IMPB have been calculated.

Te Aka Whai Ora Te Moana a Toi with District Health **Board boundaries Bay of Plenty DHB**

Figure 3 - Map of Te Moana a Toi IMPB with DHB boundaries, 2023

Table 1 shows the age breakdown of the population of Te Moana a Toi. The Māori population of Te Moana a Toi is very young, with 48% of the Māori population under the age of 25 years (compared to only 23% of the non-Māori population in the area). Over the next two decades, the Māori population is projected to grow to an estimate of 91,870 (Table 2) and to be older - by 2043, 12% of the Māori population will be over 65 years old, compared to 8% in 2023. The Māori population is projected to make up an increasing share of the IMPB population - from 25% in 2023 to 29% in 2043.

Table 1 - Population estimates by age group, Te Moana a Toi, 2023

	Māori			non-N	Total IMPB	
Age group (years)	Number	Age distribution	% of IMPB	Number	Age distribution	number
0-14	20,255	30%		30,670	15%	50,925
15-24	12,285	18%		16,810	8%	29,095
25-44	16,465	24%		50,870	25%	67,335
45-64	13,030	19%		52,935	26%	65,965
65+	5,575	8%		51,760	26%	57,335
Total	68,000	100%	25%	202,740	100%	270,740

Source: Te Whatu Ora Populations Webtool (Statistics NZ base Census 2018 base).

Table 2 - Population projections, Te Moana a Toi, 2023 to 2043

Year	Māori				non-Māori					
	Residents	% of IMPB	% 0-14 years	% 15-64 years	% 65+ years	Residents	% of IMPB	% 0-14 years	% 15-64 years	% 65+ years
2023	68,000	25%	30%	61%	8%	202,740	75%	15%	59%	26%
2028	74,250	26%	28%	62%	10%	211,340	74%	14%	58%	28%
2033	80,110	27%	27%	62%	11%	217,640	73%	13%	57%	30%
2038	86,420	28%	27%	62%	12%	221,860	72%	12%	56%	32%
2043	91,870	29%	27%	61%	12%	224,140	71%	11%	56%	33%

Source: Te Whatu Ora Populations Webtool (Statistics NZ base Census 2018 base).



The Geographic Classification for Health (GCH) is a rural-urban geographic classification composed of five categories, two urban and three rural, that reflect degrees of reducing urban influence and increasing rurality. It is applied to all of New Zealand's Statistical Areas on a scale from 'Urban 1' to 'Urban 2' based on population size, and from "Rural 1' to 'Rural 3' based on drive time to their closest major, large, medium, and small urban areas. Most Māori in Bay of Plenty DHB (75%) live in urban areas, with 24% living in rural areas compared to 93% and 6.6% for non-Māori respectively (Figure 4).

100%

80%

60%

20%

U1

U2

R1

R2

R3

■Māori ■non-Māori

Figure 4 - Population distribution by urban and rural classification, Bay of Plenty DHB, 2023

Source: Population count (Population Webtool SA2 2023); GCH (SA2 University of Otago). Note that total values may add up to more than 100% due to rounding.



Mauri ora Overall health status

3. Mauri ora - overall health status

3.1. Life Expectancy

The life expectancy at birth for Māori born in Te Moana a Toi between 2018-2022 is 78.2 years for females and 73.8 years for males (Table 3). Māori life expectancy in Te Moana a Toi is 7.5 years shorter for Māori females and 8 years shorter for Māori males, compared to non-Māori in Te Moana a Toi.

Table 3 - Life expectancy at birth, Māori and non-Māori, Te Moana a Toi, 2018 to 2022

Sex		Māori	non-Māori		Difference in years	
Sex	Years	(95% credible interval)	Years (95% credible interval)			
Female	78.2	(77.4, 79.1)	85.7	(85.3, 86.0)	-7.5	
Male	73.8	(72.9, 74.7)	81.8	(81.4, 82.2)	-8.0	

Source: Mortality data sourced from Ministry of Health. Mortality Collection, https://www.health.govt.nz/nz-health-statistics/national-collections-and-surveys/collections/mortality-collection.

Population denominator data from Statistics New Zealand, Population estimates (2022 update).

Analysed by Michael Walsh, Equity, Scientific and Technical Team, Equity Directorate, Service Improvement and Innovation, Te Whatu Ora; October 2023

In terms of the conditions which make up the life expectancy gap for Māori, this degree of information is not available at IMPB level, however analysis has been done for the four Te Whatu Ora regions of Aotearoa. In Te Manawa Taki, the region in which Te Moana a Toi is situated, for the period 2018 to 2020 the life expectancy for Māori was 75.0 years, 8.1 years lower than the non-Māori/non-Pacific population (83.1 years). Te Manawa Taki region also includes Waikato, Lakes, Taranaki, and Tairāwhiti DHBs.

Avoidable deaths include those considered *amenable* to high-quality healthcare, *preventable* through public health interventions, or both. Of the 8.1-year life expectancy gap for Māori in Te Manawa Taki, 2.7 years can be attributed to conditions that are considered both amenable and preventable followed by 1.4 years from conditions considered preventable only and 0.9 years from conditions considered amenable only. An additional 3.1 years can be attributed to conditions that are considered non avoidable ⁷.

The leading avoidable causes of death contributing to the life expectancy gap among Māori in Te Manawa Taki are lung cancer (0.9 years), coronary disease (0.8 years) and diabetes (0.6 years). A list of the top 10 conditions and their contribution to the gap are presented in Table 4. In total, these conditions contribute 4.1 years of the 8.1-year gap. These data are not able to be disaggregated by sex for Māori at a regional level because the numbers are too small.

-

⁷ By 'non-avoidable', the metric is referring to the direct causal pathway. Broader determinants of health such as income, education, housing, colonisation and institutional racism are not covered. Longer term all the 'gap' is avoidable through government, policy and intersectoral actions.

Table 4 - Decomposition of the ethnic gap in life expectancy by avoidable category - Māori compared with non-Māori/non-Pacific, 2018 to 2020, Te Manawa Taki Region

Avoidable cause	Contribution (years)
Lung cancer	0.9
Coronary disease	0.8
Diabetes	0.6
Chronic obstructive pulmonary disease	0.4
Land transport injuries	0.4
Stroke	0.3
Suicide	0.3
Valvular heart disease	0.2
Other accidental injuries	0.2
Breast cancer	0.1
Total contribution from top 10 avoidable conditions	4.1 years*

Source: Te Whatu Ora, May 2023. The Contribution of Avoidable Mortality to the Life Expectancy Gap among the Māori and Pacific population. Regional Summary.

Note: * total number provided reflects source reporting (rounding issues may apply).

3.2. Self-assessed health

In 2018, 83.3% of Māori aged 15 years and over in Te Moana a Toi reported their own health status as good, very good or excellent (Table 5), a similar percentage to Māori nationally (82.3%). A total of 16.6% of Māori in Te Moana a Toi reported their health status as fair or poor.

Table 5 - Health status reported by Māori aged 15 years and over, Te Moana a Toi, 2018

Health Status	-	Ге Moana a Toi	Aotearoa		
nealth Status	%	(95% CI)	%	(95% CI)	
Excellent	13.0	(9.8, 16.2)	15.1	(14.0, 16.2)	
Very Good	40.2	(35.6, 44.9)	36.9	(35.4, 38.3)	
Good	30.1	(25.3, 35.0)	30.3	(29.0, 31.7)	
Fair/poor	16.6	(12.9, 20.3)	17.7	(16.6, 18.8)	

Source: Te Kupenga 2018, Statistics New Zealand customised report.



3.3. Mortality

The leading causes of death for Māori in Bay of Plenty DHB in 2014-2018 were ischaemic heart disease, lung cancer, diabetes, chronic obstructive pulmonary disease (COPD) and cerebrovascular disease (Table 6), and this aligns with the leading causes of death for Māori nationally (Table 7). This differs to the leading causes of death for non-Māori in Bay of Plenty DHB, which were ischaemic heart disease, dementia, cerebrovascular disease, lung cancer and COPD in 2014-2018.

The leading causes of death for Māori females in Bay of Plenty DHB in 2014-2018 were ischaemic heart disease, lung cancer, cerebrovascular disease, COPD and breast cancer (Table 6). For Māori males, the leading causes of death in 2014-2018 were ischaemic heart disease, lung cancer, diabetes, COPD and transport accidents. Because of the small population size of a single DHB, just 1-2 deaths from a particular cause can have an impact on the ranking of leading causes. For this reason, local causes of death for Māori men and women should be interpreted together with the leading causes of death for Māori nationally (Table 7).

Table 6 - Leading causes of death for Māori, all ages, Bay of Plenty DHB, 2014 to 2018

		Mā	iori		non-N	/lāori				
Cause	Av. no. per year		e-standardised e per 100,000 (95% CI)	Av. no. per year	Age-standardised rate per 100,000 (95% CI)		J	Māori/non-Māori rate ratio (95% CI)		
Female										
Ischaemic heart disease	19	28.4	(16.2, 45.5)	98	8.3	(6.2, 10.9) 3.40	(1.95, 5.93)	20.1	
Lung cancer	18	29.3	(17.0, 46.8)	33	6.7	(3.9, 10.4) 4.36	(2.25, 8.46)	22.6	
Cerebrovascular disease	12	17.7	(8.4, 32.0)	76	7.0	(4.6, 9.8)	2.52	(1.24, 5.13)	10.7	
COPD	12	17.6	(8.5, 31.8)	37	5.2	(3.2, 7.8)	3.41	(1.63, 7.16)	12.4	
Breast cancer	10	19.1	(8.2, 36.8)	30	8.8	(4.7, 14.3) 2.17	(0.92, 5.10)	10.3	
Male										
Ischaemic heart disease	30	57.0	(37.5, 82.7)	131	20.6	(15.7, 26.1) 2.77	(1.77, 4.35)	36.4	
Lung cancer	16	32.5	(18.3, 53.1)	45	9.7	(6.2, 14.1) 3.34	(1.78, 6.28)	22.8	
Diabetes mellitus	14	26.5	(13.9, 45.4)	14	2.8	(1.0, 5.5)	9.39	(3.75, 23.49)	23.7	
COPD	8	15.2	(6.5, 29.7)	38	4.7	(3.1, 6.7)	3.25	(1.48, 7.12)	10.5	
Transport accidents	7	25.3	(10.1, 51.8)	13	11.6	(4.6, 22.3) 2.18	(0.79, 6.04)	13.7	
Total	•				•		•		•	
Ischaemic heart disease	49	41.3	(29.9, 55.5)	229	14.1	(11.5, 16.9) 2.94	(2.06, 4.19)	27.2	
Lung cancer	34	30.9	(21.1, 43.4)	78	8.1	(5.8, 10.8	3.80	(2.41, 5.99)	22.8	
Diabetes mellitus	23	20.3	(12.5, 31.1)	26	2.1	(1.0, 3.4)	9.92	(4.93, 19.95)	18.2	
COPD	20	16.2	(9.7, 25.4)	75	4.9	(3.5, 6.5)	3.33	(1.94, 5.72)	11.3	
Cerebrovascular disease	17	15.5	(8.6, 25.4)	127	7.2	(5.2, 9.6)	2.14	(1.20, 3.85)	8.3	

Source: Mortality dataset, Ministry of Health.

Notes: Ratios in **bold** show that Māori rates were significantly different from non-Māori rates in the DHB. Cerebrovascular disease includes stroke.



Table 7 - Leading causes of death for Māori, all ages, Aotearoa, 2014 to 2018

	Māori Age-standardised rate per 100,000 (95% CI)		n	on-Māori						
Cause			Age-standardised rate per 100,000 (95% CI)			ri/non-Māori ratio (95% CI)	non-Māori leading cause			
Female										
Lung cancer	29.4	(25.4, 33.9)	7.7	(7.0, 8.4)	3.84	(3.24, 4.55)	Ischaemic heart disease			
Ischaemic heart disease	24.4	(20.8, 28.3)	10.1	(9.5, 10.7)	2.42	(2.05, 2.84)	Dementia			
COPD	16.6	(13.7, 19.9)	5.3	(4.8, 5.8)	3.14	(2.55, 3.86)	Cerebrovascular disease			
Cerebrovascular disease	13.9	(11.2, 17.1)	7.7	(7.1, 8.4)	1.80	(1.44, 2.25)	COPD			
Diabetes mellitus	12.9	(10.3, 16.0)	2.7	(2.3, 3.2)	4.76	(3.64, 6.23)	Lung cancer			
Male										
Ischaemic heart disease	56.7	(50.5, 63.4)	25.3	(24.1, 26.6)	2.24	(1.98, 2.53)	Ischaemic heart disease			
Lung cancer	28.4	(24.2, 33.2)	9.1	(8.4, 9.9)	3.12	(2.61, 3.72)	Dementia			
Diabetes mellitus	19.3	(15.8, 23.4)	4.1	(3.6, 4.6)	4.76	(3.77, 6.00)	Cerebrovascular disease			
COPD	15.5	(12.5, 19.1)	6.4	(5.8, 6.9)	2.44	(1.95, 3.04)	Lung cancer			
Suicide	23.6	(18.8, 29.3)	13.0	(11.4, 14.6)	1.82	(1.42, 2.34)	COPD			
Total										
Ischaemic heart disease	39.4	(35.9, 43.1)	17.3	(16.6, 18.0)	2.27	(2.06, 2.51)	Ischaemic heart disease			
Lung cancer	29.0	(26.0, 32.2)	8.3	(7.8, 8.9)	3.48	(3.08, 3.93)	Dementia			
COPD	16.0	(13.9, 18.3)	5.7	(5.4, 6.1)	2.79	(2.40, 3.24)	Cerebrovascular disease			
Diabetes mellitus	15.9	(13.7, 18.4)	3.4	(3.0, 3.7)	4.75	(3.99, 5.67)	Lung cancer			
Cerebrovascular disease	13.4	(11.4, 15.7)	8.0	(7.5, 8.4)	1.68	(1.43, 1.99)	COPD			

Source: Mortality dataset, Ministry of Health.

Note: Ratios in **bold** show that Māori rates were significantly different from non-Māori rates. Cerebrovascular disease includes stroke. Dementia includes Alzheimer's Disease.

When looking at all deaths, the age-standardised death rate (332 deaths each year per 100,000 people) was 2.1 times higher for Māori compared to non-Māori in Bay of Plenty DHB in 2014-2018 (Table 8). This equates to an average of 159 Māori females and 176 Māori males dying each year in Bay of Plenty DHB.

Table 8 - All-cause deaths, all ages, Bay of Plenty DHB, 2014 to 2018

Māori				non-N	lāori					
Sex	Av. no. per year	Age-standardised rate per 100,000 (95% CI)		Av. no. per year		-standardised e per 100,000 (95% CI)		ori/non-Māori ratio (95% CI)	Rate difference	
Female	159	282.6	(235.5, 335.6)	800	124.6	(105.8, 144.3)	2.27	(1.80, 2.86)	158.0	
Male	176	388.3	(329.0, 454.6)	831	190.5	(166.1, 216.2)	2.04	(1.66, 2.50)	197.8	
Total	335	332.1	(294.2, 373.1)	1,631	156.7	(141.2, 172.8)	2.12	(1.82, 2.47)	175.4	

Source: Mortality dataset, Ministry of Health.

Note: Ratios in **bold** show that Māori rates were significantly different from non-Māori rates in the DHB. Average no. per year columns may not total exactly because of rounding.



The gap between Māori and non-Māori was higher for avoidable deaths (those deaths considered amenable to high-quality healthcare, preventable through public health interventions, or both) compared to all deaths in Bay of Plenty DHB (Table 9). The age-standardised potentially avoidable death rate (205 deaths each year per 100,000 people) was 2.6 times higher for Māori compared to non-Māori in Bay of Plenty DHB in 2014-2018. This equates to an average of 82 avoidable deaths each year in Māori females aged 0-74 years, and 94 in Māori males in Bay of Plenty DHB.

Table 9 - Potentially avoidable deaths, ages 0-74 years, Bay of Plenty DHB, 2014 to 2018

Māori				non-N	lāori				Rate difference	
Sex	Av. no. per year	Age-standardised rate per 100,000 (95% CI)		Av.no. per year	Age-standardised rate per 100,000 (95% CI)					
Female	82	178.9	(139.6, 225.3)	121	60.1	(44.2,	78.4)	2.98	(2.07, 4.27)	118.8
Male	94	233.5	(186.0, 288.9)	180	99.6	(78.9,	122.8)	2.34	(1.73, 3.18)	133.9
Total	176	205.2	(174.1, 240.0)	301	79.8	(66.5,	94.2)	2.57	(2.04, 3.25)	125.4

Source: Mortality dataset, Ministry of Health.

Note: Ratios in bold show that Māori rates were significantly different from non-Māori rates in the DHB.

The leading causes of potentially avoidable deaths (those deaths considered amenable to high-quality healthcare, preventable through public health interventions, or both) for Māori aged 0-74 years in Bay of Plenty DHB were ischaemic heart disease, lung cancer, diabetes, COPD and suicide (Table 10), and this aligns to the leading causes of potentially avoidable death for Māori nationally (Table 11).

The leading causes of potentially avoidable deaths for Māori females in Bay of Plenty DHB in 2014-2018 were lung cancer, ischaemic heart disease, breast cancer, cerebrovascular disease and COPD (Table 10). For Māori males, the leading causes of death in 2014-2018 were ischaemic heart disease, lung cancer, diabetes, suicide and motor vehicle accidents. Because of the small population size of a single DHB, just 1-2 deaths from a particular cause can have an impact on the ranking of leading causes. For this reason, local causes of potentially avoidable death for Māori men and women should be interpreted together with the leading causes of potentially avoidable death for Māori nationally (Table 11).

Māori aged 0-74 years in Bay of Plenty DHB in 2014-2018 had over 12.6 times higher potentially avoidable mortality from diabetes compared to non-Māori, 4.3 times higher potentially avoidable mortality for lung cancer, 4.1 times higher potentially avoidable mortality for COPD and 4.0 times higher potentially avoidable mortality for ischaemic heart disease (Table 10).



Table 10 - Leading causes of potentially avoidable mortality, ages 0-74 years, Bay of Plenty DHB, 2014 to 2018

		Māori				lāori				
Cause	Av. no. per year	rate per 100 000 rate per 100 000		per 100,000	Māo rate	Rate difference				
Female										
Lung cancer	14	25.8	(14.1, 43.2)	16	5.3	(2.7, 9.2)	4.85	(2.24, 10.48)	20.5	
Ischaemic heart disease	10	19.6	(9.1, 36.4)	11	3.0	(1.4, 5.6)	6.45	(2.61, 15.93)	16.6	
Breast cancer	8	17.0	(6.6, 35.0)	17	7.8	(3.7, 13.6)	2.19	(0.85, 5.68)	9.2	
Cerebrovascular disease	7	13.3	(5.1, 27.8)	8	2.7	(0.9, 5.9)	4.88	(1.59, 14.98)	10.6	
COPD	7	12.8	(4.9, 26.7)	13	3.4	(1.6, 6.1)	3.75	(1.42, 9.94)	9.4	
Male										
Ischaemic heart disease	19	40.5	(23.5, 64.5)	33	11.9	(7.5, 17.5)	3.41	(1.83, 6.36)	28.6	
Lung cancer	14	29.6	(15.9, 49.9)	22	7.5	(4.2, 12.0)	3.96	(1.92, 8.20)	22.1	
Diabetes	11	23.5	(11.5, 42.2)	6	2.1	(0.5, 5.1)	11.39	(3.62, 35.85)	21.4	
Suicide and self- inflicted injuries	7	24.2	(9.3, 50.8)	15	16.6	(8.1, 29.0)	1.46	(0.56, 3.83)	7.6	
Motor vehicle accidents	6	22.3	(8.2, 47.8)	8	9.9	(3.5, 20.7)	2.25	(0.74, 6.86)	12.4	
Total										
Ischaemic heart disease	29	29.4	(19.3, 42.9)	44	7.2	(4.9, 10.1)	4.06	(2.43, 6.79)	22.2	
Lung cancer	29	27.5	(18.2, 39.9)	39	6.4	(4.2, 9.1)	4.34	(2.56, 7.35)	21.1	
Diabetes	17	17.1	(9.7, 27.6)	8	1.4	(0.5, 2.9)	12.56	(4.93, 32.01)	15.7	
COPD	12	11.1	(5.5, 19.8)	21	2.7	(1.6, 4.3)	4.08	(1.92, 8.68)	8.4	
Suicide and self- inflicted injuries	11	18.5	(9.0, 33.7)	18	9.9	(5.2, 16.6)	1.87	(0.83, 4.20)	8.6	

Source: Mortality dataset, Ministry of Health.

Notes: Ratios in **bold** show that Māori rates were significantly different from non-Māori rates in the DHB. Cerebrovascular disease includes stroke.



Table 11 - Leading causes of potentially avoidable mortality, ages 0-74 years, Aotearoa, 2014 to 2018

	Māori		n	on-Māori						
Cause	rate	standardised per 100,000 (95% CI)	rate	standardised per 100,000 (95% CI)	Māori/non-Māori rate ratio (95% CI)		non-Māori Ieading cause			
Female										
Lung cancer	24.6	(20.8, 28.9)	6.0	(5.3, 6.7)	4.11	(3.38, 5.00)	Breast cancer			
Ischaemic heart disease	14.5	(11.5, 17.9)	3.9	(3.4, 4.5)	3.67	(2.85, 4.74)	Lung cancer			
COPD	11.2	(8.7, 14.1)	3.1	(2.7, 3.6)	3.59	(2.72, 4.74)	Ischaemic heart disease			
Breast cancer	11.7	(8.9, 15.1)	8.1	(7.2, 9.1)	1.45	(1.09, 1.92)	Colorectal cancer			
Diabetes	9.7	(7.3, 12.6)	1.7	(1.4, 2.2)	5.56	(3.91, 7.91)	COPD			
Male										
Ischaemic heart disease	42.1	(36.7, 48.1)	15.5	(14.4, 16.7)	2.71	(2.33, 3.16)	Ischaemic heart disease			
Lung cancer	24.0	(20.1, 28.5)	6.7	(6.0, 7.5)	3.59	(2.93, 4.40)	Lung cancer			
Suicide and self-inflicted injuries	23.8	(18.9, 29.5)	12.9	(11.4, 14.6)	1.84	(1.43, 2.36)	Suicide and self-inflicted injuries			
Diabetes	15.5	(12.3, 19.3)	2.8	(2.3, 3.3)	5.64	(4.24, 7.51)	Colorectal cancer			
Motor vehicle accidents	16.1	(12.2, 20.7)	7.0	(5.8, 8.4)	2.29	(1.68, 3.13)	Cerebrovascular disease			
Total		,								
Ischaemic heart disease	27.6	(24.5, 30.9)	9.6	(9.0, 10.2)	2.88	(2.52, 3.28)	Ischaemic heart disease			
Lung cancer	24.3	(21.6, 27.4)	6.3	(5.8, 6.8)	3.85	(3.34, 4.43)	Lung cancer			
Diabetes	12.4	(10.4, 14.7)	2.2	(1.9, 2.6)	5.58	(4.47, 6.96)	Colorectal cancer			
Suicide and self-inflicted injuries	16.9	(14.0, 20.2)	8.6	(7.7, 9.6)	1.96	(1.59, 2.41)	Suicide and self-inflicted injuries			
COPD	10.4	(8.6, 12.4)	3.2	(2.8, 3.5)	3.30	(2.68, 4.05)	COPD			

Source: Mortality dataset, Ministry of Health.

Note: Ratios in **bold** show that Māori rates were significantly different from non-Māori rates. Cerebrovascular disease includes stroke.





4. Whānau ora - Healthy families

Māori models of health encompass cultural vitality and whānau wellbeing. Indicators of these dimensions of health specific for Māori in each IMPB are included in these profiles, sourced from Te Kupenga 2018, the Māori Social Survey conducted in 2018 by StatsNZ. In 2018, this was a survey of almost 8,500 adults (aged 15 years and over) of Māori ethnicity and/or descent. Further information on Te Kupenga can be found here. Data from Te Kupenga are presented for Māori only.

Based on a scale where 0 is doing extremely badly and 10 is doing extremely well (Table 12), most Māori (76.7%) in Te Moana a Toi reported their whānau was doing well (7/10 or greater), compared to 73.6% of Māori nationally. Just under a quarter of Māori (23.2%) in Te Moana a Toi reported that their whānau was not doing well (6/10 or less).

Table 12 - Whānau well-being reported by Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018

How the whēney is doing	Т	e Moana a Toi	Aotearoa		
How the whānau is doing	%	(95% CI)	%	(95% CI)	
(10 out of 10)	16.9	(13.0, 20.9)	12.9	(12.1, 13.7)	
(9 out of 10)	13.3 *	(9.1, 17.6)	12.8	(11.9, 13.6)	
(8 out of 10)	25.9	(21.4, 30.4)	24.4	(23.3, 25.6)	
(7 out of 10)	20.6	(16.1, 25.1)	23.5	(22.5, 24.6)	
(0-6 out of 10)	23.2	(18.8, 27.6)	26.4	(25.2, 27.6)	

Source: Te Kupenga 2018, Statistics New Zealand customised report.

Note: An asterisk (*) shows the sampling error is 30% or more but less than 50%.

When thinking about who made up the whānau, about 22% of Māori in Te Moana a Toi included "close friends or others" (Table 13).

Table 13 - Whānau composition reported by Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018

Whānau dagarintian	Te	Moana a Toi	Aotearoa		
Whānau description	% (95% CI)		%	(95% CI)	
Size of whānau					
10 or less	47.6	(41.9, 53.4)	52.1	(50.6, 53.6)	
11 to 20	23.2	(18.7, 27.7)	24.2	(23.0, 25.4)	
More than 20	29.2	(24.1, 34.3)	23.7	(22.3, 25.0)	
Groups included in whānau					
Parents, partner, children, brothers and sisters	97.6	(95.5, 99.6)	97.4	(97.0, 97.8)	
Grandparents, grandchildren	45.0	(38.6, 51.5)	39.0	(37.5, 40.5)	
Aunts and uncles, cousins, nephews and nieces, other in-laws	48.6	(42.6, 54.7)	48.6	(47.1, 50.2)	
Close friends, others	22.0 *	(15.1, 29.0)	22.6	(21.3, 23.8)	

Source: Te Kupenga 2018, Statistics New Zealand customised report.

Notes: An asterisk (*) shows the sampling error is 30% or more but less than 50%.

⁸ https://www.stats.govt.nz/information-releases/te-kupenga-2018-final-english

Most Māori (77.3) in Te Moana a Toi reported it was easy or very easy to get support in times of need. Fewer Māori (64.4%) reported it was easy or very easy to get help with Māori cultural practices, such as going to a tangi, speaking at a hui or blessing a tanga (Table 14).

Table 14 - Access to whānau support, Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018

How convict to not halp	Te Moan	a a Toi	Aotearoa				
How easy is it to get help	% (95% CI)		%	(95% CI)			
Support in times of need							
Easy, very easy	77.3	(73.3, 81.2)	76.1	(74.9, 77.3)			
Sometimes easy, sometimes hard	13.0	(9.5, 16.5)	16.4	(15.5, 17.4)			
Hard, very hard	9.7	(7.1, 12.4)	7.5	(6.7, 8.3)			
Help with Māori cultural practices such	as going to a tangi, sp	eaking at a hui, or b	lessing a ta	ionga			
Easy, very easy	64.4	(59.0, 69.8)	59.0	(57.7, 60.3)			
Sometimes easy, sometimes hard	14.7	(10.4, 18.9)	18.9	(17.9, 19.9)			
Hard, very hard	15.9	(11.3, 20.6)	18.1	(17.0, 19.2)			

Source: Te Kupenga 2018, Statistics New Zealand customised report.

Being involved in Māori culture was very/quite important to 49.1% of Māori in Te Moana a Toi, and spirituality was very/quite important to 55.7% of Māori in Te Moana a Toi (Table 15). Only 10.9% of Māori respondents in Te Moana a Toi reported that being involved in Māori culture was not at all important to them.

Table 15 - Importance of Māori culture and spirituality, Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018

	-	Ге Moana a Toi		Aotearoa						
	%	(95% CI)	%	(95% CI)						
Importance of being involved in Māori culture										
Very important	25.5	(21.4, 29.6)	22.1	(21.1, 23.1)						
Quite important	23.6	(19.6, 27.6)	23.2	(22.1, 24.3)						
Somewhat	23.6	(19.9, 27.3)	25.8	(24.7, 26.9)						
A little important	16.4	(11.3, 21.6)	18.3	(17.1, 19.5)						
Not at all important	10.9	(7.4, 14.4)	10.6	(9.7, 11.6)						
Importance of spirituality										
Very important	36.7	(31.7, 41.6)	30.7	(29.5, 31.9)						
Quite important	19.0	(15.4, 22.5)	18.0	(16.9, 19.0)						
Somewhat	15.2	(11.6, 18.9)	16.8	(15.9, 17.8)						
A little important	11.6	(8.6, 14.6)	15.3	(14.3, 16.2)						
Not at all important	17.5	(13.1, 22.0)	19.2	(18.1, 20.4)						

Source: Te Kupenga 2018, Statistics New Zealand customised report.



In Te Moana a Toi in 2018, just over a quarter of Māori (27.7%) used te reo Māori regularly in the home, compared to 18.4% of Māori nationally (Table 16). Small numbers in the Te Kupenga survey sample make it unreliable to determine the percentage of homes where te reo Māori is the main language in Te Moana a Toi.

Table 16 - Use of te reo Māori in the home, Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018

Language spoken at home	Т	e Moana a Toi	Aotearoa		
	%	(95% CI)	%	(95% CI)	
Māori is main language	S	(NA, NA)	1.8	(1.3, 2.2)	
Māori is used regularly	used regularly 27.7		18.4	(17.3, 19.5)	

Source: Te Kupenga 2018, Statistics New Zealand customised report.

Notes: NA = Not Available, S = suppressed: number too small for reliable estimate.

In 2018, almost all Māori in Te Moana a Toi (98.7%) had been to a marae, with a majority (63.9%) having been in the last 12 months (Table 17). Of those who had ever been to a marae, and who knew their ancestral marae, 85.5% had been to an ancestral marae at some time, over a half (54.1%) had been in the last 12 months and 56.9% would like to go more often.

Table 17 - Access to marae, Māori aged 15 years and over, Te Moana a Toi and Aotearoa, 2018

Been to marae	Т	e Moana a Toi	Aotearoa		
Deen to marae	%	(95% CI)	%	(95% CI)	
At some time	98.7	(97.1, 100.2)	96.6	(96.0, 97.1)	
In previous 12 months [1]	63.9	(58.2, 69.5)	51.8	(50.6, 53.1)	
Ancestral marae at some time [1][2]	85.5	(81.1, 89.8)	84.3	(82.9, 85.6)	
Ancestral marae in previous 12 months [1][2]	54.1	(48.2, 59.9)	44.3	(42.6, 45.9)	
Like to go to ancestral marae more often [1][2]	56.9	(50.7, 63.0)	63.6	(62.1, 65.1)	

Source: Te Kupenga 2018, Statistics New Zealand customised report.

Notes: [1] Those who had been to a marae at some time. [2] Includes only those who knew their ancestral marae.

18.2% of Māori aged 15 years and over in Te Moana a Toi had taken part in traditional healing or massage in the past 12 months in 2018, compared to 12.3% of Māori nationally (Table 18).

Table 18 - Māori aged 15 years and over who took part in traditional healing or massage in last 12 months, Te Moana a Toi and Aotearoa, 2018

	Te Moana a Toi		Aotearoa
%	(95% CI)	%	(95% CI)
18.2	(14.2, 22.1)	12.3	(11.4, 13.2)

Source: Te Kupenga 2018, Statistics New Zealand customised report.





5. Wai ora – Healthy environments

This section focuses on key aspects of social and physical environments that influence health and well-being. Information in this section comes from Māori and non-Māori individuals responding to the NZ Census 2018, or Māori respondents in the 2018 Te Kupenga survey. Because of data availability at the time of writing, NZ Census2018, NZDep2018 and PHO enrolment data are presented for the Bay of Plenty DHB geographical area, whereas Te Kupenga survey data is presented for the Te Moana a Toi IMPB geographic area. The data quality and degree of certainty for Māori is not the same for all variables from the NZ Census 2018. Please see the technical appendix at the end of this report, for further details about how geographic areas were defined for each data source, and for more information on how to interpret variables from the NZ Census 2018.

5.1. Education

In 2018, 66.3% of Māori aged over 20 years in Bay of Plenty DHB had achieved a Level 2 Certificate or higher, compared to 79.3% for non-Māori (Table 19).

Table 19 - Adults aged 20 years and over with a Level 2 Certificate or higher, Bay of Plenty DHB, 2018

		Māo	ri		non-N	lāori	Mā	ori/non-Māori	Difference
Year	Number	%	(95% CI)	Number	%	(95% CI)	rate	ratio (95% CI)	in percentage
2018	21,402	66.3	(65.4, 67.2)	91,686	79.3	(78.7, 80.0)	0.84 (0.83, 0.84)		-13.0

Source: 2018 Census, Statistics New Zealand.

Notes: Percentages are age-standardised to the 2001 Māori population. Ratios in **bold** show a statistically significant difference between Māori and non-Māori.

5.2. Work

In 2018, 47.4% of Māori aged 15 years and over in Bay of Plenty DHB were employed full time, and 16.4% were employed part time (Table 20). In 2018, 9.5% of Māori in Bay of Plenty DHB were unemployed, over twice the rate of non-Māori, and Māori were 1.2 times more likely than non-Māori to not be in the labour force.

Table 20 - Labour force status, 15 years and over, Bay of Plenty DHB, 2018

Labour force	Māori				non-Māori			ori/non-Māori	Difference
status	Number	%	(95% CI)	Number	%	(95% CI)	rate ratio (95% CI)		percentage
Employed full- time	19,230	47.4	(46.7, 48.1)	69,507	55.5	(55.0, 55.9)	0.85	(0.84, 0.86)	-8.1
Employed part- time	6,660	16.4	(15.9, 16.8)	22,764	17.7	(17.5, 18.0)	0.92	(0.90, 0.94)	-1.4
Unemployed	3,573	9.5	(9.1, 9.8)	4,053	4.2	(4.0, 4.3)	2.26	(2.16, 2.36)	5.3
Not in the labour force	12,567	26.9	(26.4, 27.4)	52,875	22.6	(22.3, 22.9)	1.19	(1.17, 1.21)	4.3

Source: 2018 Census, Statistics New Zealand

Notes: Percentages are age-standardised to the 2001 Māori population. Ratios in **bold** show a statistically significant difference between Māori and non-Māori. Employed part-time includes people working 1 hour per week or more. Employed full-time includes people who usually work 30 or more hours per week. Unemployed people are without a paid job, available for work and actively seeking work. People not in the labour force includes people in the working age population who are neither employed nor unemployed.

In 2018, the main employers of Māori women in Bay of Plenty DHB were health care and social assistance (17.2%); education and training (15.0%); retail trade (10.8%); accommodation and food services (9.3%); and administrative and support services (9.0%) (Table 21). For Māori men, the leading industries were construction (19.3%); manufacturing (15.1%); agriculture, forestry and fishing (13.3%); transport, postal and warehousing (8.8%); and administrative and support services (7.8%).

Table 21 - Leading industries in which Māori were employed, Bay of Plenty DHB, 2018

ANZSIC Industry		Māori		non-Māori			
	Number	%	Rank	Number	%	Rank	
Females							
Health Care and Social Assistance	2,181	17.2%	1	8,232	18.8%	1	
Education and Training	1,902	15.0%	2	5,187	11.8%	2	
Retail Trade	1368	10.8%	3	4,782	10.9%	3	
Accommodation and Food Services	1176	9.3%	4	2,871	6.6%	5	
Administrative and Support Services	1143	9.0%	5	2,298	5.2%	8	
Males							
Construction	2,550	19.3%	1	8,310	17.1%	1	
Manufacturing	1,995	15.1%	2	6,147	12.7%	2	
Agriculture, Forestry and Fishing	1,758	13.3%	3	5,295	10.9%	3	
Transport, Postal and Warehousing	1,170	8.8%	4	3,126	6.4%	6	
Administrative and Support Services	1,026	7.8%	5	2,418	5.0%	8	

Source: 2018 Census, Statistics New Zealand.

Note: Australian and New Zealand Standard Industrial Classification (ANZSIC).



In terms of the type of work Māori perform within those industries (Table 22), for employed Māori women in Bay of Plenty DHB, the leading occupational groupings were professionals (21.5%); community and personal service workers (18.3%); labourers (16.2%); and clerical and administrative workers (14.6%). Māori men were most likely to be employed as labourers (26.5%); technicians and trades workers (17.5%); machinery operators and drivers (17.3%); and managers (15.2%).

Table 22 - Leading occupations in which Māori were employed, Bay of Plenty DHB, 2018

ANZSCO Occupation		Māori		non-Māori			
	Number	%	Rank	Number	%	Rank	
Females							
Professionals	2,718	21.5%	1	11,673	26.7%	1	
Community and Personal Service Workers	2,316	18.3%	2	5,943	13.6%	4	
Labourers	2,049	16.2%	3	3,450	7.9%	6	
Clerical and Administrative Workers	1,851	14.6%	4	8,373	19.1%	2	
Sales Workers	1,470	11.6%	5	4,953	11.3%	5	
Managers	1,428	11.3%	6	6,552	15.0%	3	
Technicians and Trades Workers	531	4.2%	7	2,367	5.4%	7	
Machinery Operators and Drivers	282	2.2%	8	474	1.1%	8	
Males							
Labourers	3,513	26.5%	1	6,675	13.8%	4	
Technicians and Trades Workers	2,316	17.5%	2	9,684	20.0%	2	
Machinery Operators and Drivers	2,289	17.3%	3	4,602	9.5%	5	
Managers	2,016	15.2%	4	12,369	25.5%	1	
Professionals	1,410	10.7%	5	7,971	16.4%	3	
Community and Personal Service Workers	681	5.1%	6	1,977	4.1%	7	
Sales Workers	624	4.7%	7	3,444	7.1%	6	
Clerical and Administrative Workers	387	2.9%	8	1,779	3.7%	8	

Source: 2018 Census, Statistics New Zealand.

Note: Australian and New Zealand Standard Classification of Occupations (ANZSCO), major grouping.



Unpaid work is very common, with 89.5% of Māori aged over 15 years in Bay of Plenty DHB in 2018 reporting they performed unpaid work (Table 23). Māori in Bay of Plenty DHB were significantly more likely than non-Māori to participate in unpaid work looking after a disabled or ill household (2.0 times) or non-household (1.5 times) member.

Table 23 - Unpaid work, 15 years and over, Bay of Plenty DHB, 2018

	Māori		non-Māori		Māo	ri/non-Māori	Difference
Unpaid work	Number	%	Number	%	rate	ratio (95% CI)	in percentage
Any unpaid work	24,735	89.5	113,382	89.5	1.00	(1.00, 1.00)	0.0
Looking after disabled/ill household member	3,726	13.5	8,637	6.8	1.98	(1.91, 2.05)	6.7
Looking after disabled/ill non-household member	3,498	12.7	10,956	8.6	1.46	(1.41, 1.52)	4.0

Source: 2018 Census, Statistics New Zealand.

Notes: Percentages are NOT age-standardised due to not having detailed age-group data available.


Ratios in **bold** show a statistically significant difference between Māori and non-Māori.

5.3. Income and standard of living

NZDep2018 is a small-area-based measure of neighbourhood deprivation, by looking at the comparative socio-economic positions of small geographic areas and assigning them decile numbers from 1 (least deprived) to 10 (most deprived). The index is based on 9 socio-economic variables from the 2018 Census (Atkinson, Salmond et al. 2019). It describes the general socio-economic deprivation of an area. An area's decile score does not necessarily mean all individuals living in that area experience an equivalent level of deprivation.

In Bay of Plenty DHB, 47% of Māori lived in the two most deprived deciles in 2018, compared to 13% for non-Māori (Figure 5). A total of 7% of Māori in Bay of Plenty DHB lived in the two least deprived deciles in 2018, compared to 20% of non-Māori in Bay of Plenty DHB.

Figure 5 - NZDep2018 distribution of Māori and non-Māori by decile, Bay of Plenty DHB, 2018



Source: Deprivation decile for estimated resident population (ERP), former DHB areas, prioritised ethnicity, provided by Stats NZ for Te Whatu Ora. Deprivation is derived according to the neighbourhood where the individual lives, based on University of Otago's NZDep2018 Socio-economic Deprivation Indices.



In 2018, 11.4% of Māori aged over 15 years in Te Moana a Toi reported often postponing or putting off a doctor's visit, 6.0% often went without fresh fruit and vegetables, and 10.9% often put up with feeling cold, because of cost (Table 24).

Table 24 - Unmet needs reported by Māori aged 15 years and over to keep costs down in the last 12 months, Te Moana a Toi and Aotearoa, 2018

Actions taken a lot to keep costs down	Те	Moana a Toi	Aotearoa		
Actions taken a lot to keep costs down	%	(95% CI)	%	(95% CI)	
Put up with feeling the cold	10.9 *	(7.3, 14.5)	9.9	(9.1, 10.7)	
Go without fresh fruit and vegetables	6.0 *	(3.4, 8.6)	6.2	(5.6, 6.9)	
Postpone or put off visits to the doctor	11.4	(8.6, 14.2)	9.7	(8.8, 10.6)	

Source: Te Kupenga 2018, Statistics New Zealand customised report.

Notes: An asterisk (*) shows the sampling error is 30% or more but less than 50%. Participants were asked if they did any of these "a lot", "a little" or "not at all" to keep costs down. Only those who answered "a lot" are shown here.

Māori in Bay of Plenty DHB are significantly more likely than non-Māori (1.5 times) to receive an income of \$20,000 or less (Table 25). This equates to 37.9% of Māori aged 20 years and over lived on an income of \$20,000 or less compared to 25.8% of non-Māori in 2018.

Table 25 - People 20 years and over whose total annual personal income in \$20,000 or less, Bay of Plenty DHB, 2018

	Māori			non-Māori			Māo	ri/non-Māori	Difference
Measure	Number	%	(95% CI)	Number	%	(95% CI)		atio (95% CI)	percentage
Total income \$20,000 or less	13,911	37.9	(37.3, 38.6)	39,216	25.8	(25.4, 26.1)	1.47	(1.45, 1.50)	12.1

Source: 2018 Census, Statistics New Zealand.

Note: Percentages are age-standardised to the 2001 Māori population. Ratios in **bold** show a statistically significant difference between Māori and non-Māori.

Māori in Bay of Plenty DHB are 3.4 times more likely than non-Māori to be without access to a motor vehicle (Table 26). This equates to 4.8% of Māori (2,307 people) living in Bay of Plenty DHB with no access to a motor vehicle compared to 1.4% of non-Māori (3,639 people) in 2018.

Table 26 - People with no access to a motor vehicle, Bay of Plenty DHB, 2018

		Māc	ori		non-M	lāori	Māo	ri/non-Māori	Difference
Measure	Number % (95% CI)		Number	%	(95% CI)	rate ratio (95% CI) percentag			
2018	2,307	4.8	(4.6, 5.0)	3,639	1.4	(1.3, 1.5)	3.36	(3.20, 3.54)	3.4

Source: 2018 Census, Statistics New Zealand.

Notes: Percentages are age-standardised to the 2001 Māori population. Ratios in **bold** show a statistically significant difference between Māori and non-Māori.



Māori in Bay of Plenty DHB are also 2.9 times more likely than non-Māori to have no access to telecommunications (Table 27). This equates to 2.0% of Māori (933 people) who had no access to any form of telecommunications (a functional cellphone, telephone, or the Internet) compared to 0.7% of non-Māori (951 people) in 2018.

Table 27 - People with no access to telecommunications, Bay of Plenty DHB, 2018

		Mā	iori		non-M	āori	Māo	ri/non-Māori	Difference
Measure	Number	%	(95% CI)	Number	%	(95% CI)		ratio (95% CI)	in percentage
2018	933 2.0 (1.8, 2.1)		951	0.7	(0.6, 0.7)	2.88	(2.64, 3.15)	1.3	

Source: 2018 Census, Statistics New Zealand.

Notes: Percentages are age-standardised to the 2001 Māori population. Ratios in **bold** show a statistically significant difference between Māori and non-Māori.

5.4. Housing

Māori in Bay of Plenty DHB are less likely than non-Māori to own their home (Table 28). In 2018, 65.4% of Māori aged 20 years and over in Bay of Plenty DHB lived in a home they did not own/partly own or hold in a family trust compared to 49.9% of non-Māori.

Table 28 - Housing tenure, 20 years and over, Bay of Plenty DHB, 2018

		Mād	ori		non-N	lāori	Māc	ri/non-Māori	Difference
Housing tenure	Number	%	(95% CI)	Number	%	(95% CI)	rate	ratio (95% CI)	percentage
Owned or partly owned	8,595	28.9	(28.2, 29.6)	59,739	39.1	(38.7, 39.6)	0.74	(0.73, 0.75)	-10.2
Held in a family trust	1,764	5.7	(5.4, 6.0)	20,319	10.9	(10.7, 11.1)	0.52	(0.50, 0.55)	-5.2
Not owned; not held in a family trust	14,409	65.4	(64.2, 66.5)	41,097	49.9	(49.4, 50.5)	1.31	(1.29, 1.33)	15.4

Source: 2018 Census, Statistics New Zealand.

Note: Percentages are age-standardised to the 2001 Māori population. Ratios in **bold** show a statistically significant difference between Māori and non-Māori.

Living in an overcrowded home was 3.0 times more common for Māori than non-Māori in Bay of Plenty DHB in 2018 (Table 29). In the 2018 Census, 24.8% of Māori (11,487 people) in Bay of Plenty DHB lived in overcrowded homes compared to 8.2% of non-Māori (9,246 people).

Table 29 - People living in crowded households (requiring at least one more bedroom), Bay of Plenty DHB, 2018

		Māc	ori		non-M	āori	Māo	ri/non-Māori	Difference
Measure	Number	%	(95% CI)	Number	%	(95% CI)	rate r	atio (95% CI)	in percentage
Household crowding	11,487	24.8	(24.3, 25.3)	9,246	8.2	(8.0, 8.4)	3.03	(2.95, 3.10)	16.6

Source: 2018 Census, Statistics New Zealand.

Notes: Percentages are age-standardised to the 2001 Māori population. Ratios in **bold** show a statistically significant difference between Māori and non-Māori.



In 2018, 42.5% of Māori in Bay of Plenty DHB lived in a home that was sometimes or always damp, and 34.7% of Māori lived in a house with mould (Table 30). Māori in Bay of Plenty DHB were 2.0 times more likely than non-Māori to live in a damp home and to live in a mouldy home.

Table 30 - People experiencing housing quality issues sometimes or always, Bay of Plenty DHB, 2018

Housing		Māc	ori		non-M	lāori	Māo	ri/non-Māori	Difference
quality issues	Number % (95% CI)		Number	%	(95% CI)		ratio (95% CI)	in percentage	
Dampness	17,991	42.5	(41.9, 43.1)	26,016	21.2	(20.9, 21.5)	2.00	(1.97, 2.03)	21.3
Mould	14,850	34.7 (34.2, 35.3)		21,870	17.6	(17.3, 17.9)	1.97	(1.94, 2.01)	17.1

Source: 2018 Census, Statistics New Zealand.

Notes: Percentages are age-standardised to the 2001 Māori population. Ratios in **bold** show a statistically significant difference between Māori and non-Māori. Dampness indicator shows % people who stated their house experienced dampness sometimes or always. Mould indicator shows % people who stated their house experienced mould (of approximately A4-size or larger) sometimes or always.

Māori in Bay of Plenty DHB were also 1.3 times as likely as non-Māori to live in homes without any source of heating in 2018 (Table 31). This equates to 4.8% of Māori (2,145 people) in Bay of Plenty DHB who were without heating compared to 3.7% of non-Māori (4,530 people) in 2018.

Table 31 - People living in households where there is no source of heating, Bay of Plenty DHB, 2018

		Māo	ri		non-M	āori	Māo	ri/non-Māori	Difference
Measure	Number	%	(95% CI)	Number	%	(95% CI)		ratio (95% CI)	percentage
No source of heating	2,145	4.8	(4.6, 5.1)	4,530	3.7	(3.6, 3.8)	1.30	(1.24, 1.37)	1.1

Source: 2018 Census, Statistics New Zealand.

Note: Percentages are age-standardised to the 2001 Māori population. Ratios in **bold** show a statistically significant difference between Māori and non-Māori.



5.5. Primary Care Enrolment


In October 2023, 15.2% of Māori in Bay of Plenty DHB were not enrolled with primary health care, compared to 4% for non-Māori (Table 32). Nationally, 16.2% of Māori were not enrolled with primary health care, compared to 1.3% of non-Māori in October 2023. One partial explanation for the lower enrolment for Māori may be related to poor ethnicity data quality - this primary care enrolment data uses the ethnicity recorded in a person's National Health Index (NHI) record, and previous research has found that compared to the ethnicity that people report in the Census, the NHI undercounts Māori by 15.7%, with higher undercounts for Māori men (Harris, Paine et al. 2022). The poor ethnicity data quality makes it difficult to assess how many Māori in Bay of Plenty DHB are actually missing out on being enrolled with primary health care, and how many are actually enrolled but misclassified with a non-Māori ethnicity. It is likely that both of these factors make a contribution to the inequity in primary care enrolment data.

Table 32 - People enrolled with primary care, Bay of Plenty DHB, October 2023

		Mā	ori		non-N	lāori	Māor	i/non-Māori	Difference
Year	Number	%	(95% CI)	Number	%	(95% CI)	1111111	atio (95% CI)	in percentage
2023	62,434	434 84.8 (84.2, 85.5)		196,779	96	(95.6, 96.5)	0.88	(0.88, 0.89)	-11.2

Source: Te Whatu Ora Primary Care Enrolment data; denominator is 2023 ERP from Te Whatu Ora Population Web Tool Notes: *Percentages are crude (not age-standardised)*. Ratios in **bold** show a statistically significant difference between Māori and non-Māori.





Appendix 1: IMPB Māori population projections

Table 33 - Māori population projections, single year, Bay of Plenty DHB, by 5-year age band, 2018 to 2043

Age	Female	Male	Total									
Groups		2018			2019			2020			2021	
00-04	3,240	3,410	6,650	3,270	3,440	6,720	3,350	3,470	6,810	3,380	3,560	6,940
05-09	3,410	3,590	7,000	3,460	3,600	7,060	3,470	3,700	7,170	3,560	3,700	7,260
10-14	3,320	3,490	6,810	3,410	3,600	7,010	3,500	3,710	7,220	3,590	3,880	7,470
15-19	2,820	2,960	5,780	2,950	3,170	6,120	3,110	3,270	6,380	3,210	3,340	6,550
20-24	2,260	2,400	4,670	2,300	2,440	4,740	2,390	2,590	4,980	2,560	2,790	5,350
25-29	2,330	2,280	4,610	2,350	2,340	4,690	2,350	2,330	4,690	2,370	2,340	4,710
30-34	2,060	1,960	4,020	2,160	2,000	4,160	2,300	2,150	4,440	2,390	2,250	4,650
35-39	1,950	1,800	3,750	1,960	1,860	3,830	2,020	1,890	3,910	2,100	1,970	4,070
40-44	1,840	1,780	3,620	1,890	1,780	3,670	1,920	1,840	3,750	1,950	1,830	3,780
45-49	1,920	1,860	3,780	1,940	1,840	3,780	1,950	1,830	3,780	1,990	1,850	3,830
50-54	1,830	1,640	3,480	1,840	1,690	3,530	1,850	1,730	3,580	1,870	1,780	3,650
55-59	1,820	1,610	3,430	1,850	1,620	3,470	1,900	1,650	3,540	1,900	1,640	3,540
60-64	1,400	1,210	2,620	1,450	1,260	2,710	1,550	1,330	2,870	1,610	1,430	3,030
65-69	1,040	970	2,020	1,130	1,020	2,150	1,170	1,040	2,210	1,260	1,060	2,320
70-74	670	590	1,260	710	620	1,330	770	710	1,480	840	780	1,630
75-79	450	390	840	460	420	870	480	410	890	480	410	890
80-84	260	210	470	250	210	460	270	210	480	300	220	530
85+	200	100	300	220	100	320	210	130	340	220	130	340
All Ages	32,800	32,300	65,100	33,600	33,000	66,600	34,600	34,000	68,500	35,600	35,000	70,500



Age	Female	Male	Total									
Groups		2022			2023			2024			2025	
00-04	3,440	3,590	7,030	3,450	3,620	7,060	3,470	3,660	7,130	3,500	3,690	7,190
05-09	3,560	3,740	7,300	3,550	3,780	7,340	3,580	3,810	7,380	3,620	3,790	7,410
10-14	3,620	3,910	7,530	3,730	3,940	7,680	3,770	3,950	7,720	3,750	4,000	7,750
15-19	3,330	3,480	6,810	3,440	3,670	7,100	3,530	3,770	7,310	3,600	3,860	7,470
20-24	2,750	2,940	5,690	2,910	3,000	5,910	3,020	3,190	6,220	3,150	3,270	6,420
25-29	2,380	2,360	4,740	2,290	2,400	4,690	2,310	2,420	4,730	2,370	2,540	4,900
30-34	2,450	2,340	4,790	2,540	2,400	4,940	2,540	2,440	4,980	2,520	2,400	4,920
35-39	2,150	2,040	4,180	2,240	2,080	4,320	2,330	2,120	4,450	2,450	2,240	4,690
40-44	2,030	1,860	3,890	2,080	1,880	3,960	2,080	1,940	4,020	2,120	1,940	4,060
45-49	1,920	1,820	3,750	1,900	1,830	3,730	1,940	1,820	3,760	1,950	1,860	3,810
50-54	1,920	1,810	3,740	1,950	1,870	3,820	1,960	1,850	3,810	1,950	1,820	3,770
55-59	1,890	1,620	3,510	1,840	1,620	3,470	1,840	1,670	3,510	1,840	1,690	3,530
60-64	1,670	1,520	3,190	1,790	1,550	3,350	1,830	1,560	3,400	1,860	1,590	3,450
65-69	1,320	1,110	2,420	1,360	1,140	2,490	1,400	1,180	2,580	1,480	1,230	2,710
70-74	900	820	1,720	940	850	1,800	1,020	900	1,930	1,060	920	1,980
75-79	510	430	950	550	470	1,020	590	490	1,080	640	570	1,200
80-84	310	240	550	330	270	590	330	290	620	350	280	630
85+	230	140	360	230	130	360	230	130	360	230	150	390
All Ages	36,400	35,800	72,100	37,100	36,500	73,600	37,800	37,200	75,000	38,400	37,800	76,300



Age	Female	Male	Total									
Groups		2026			2027			2028			2029	
00-04	3,530	3,720	7,250	3,570	3,770	7,340	3,620	3,820	7,440	3,660	3,850	7,510
05-09	3,610	3,850	7,460	3,650	3,860	7,510	3,650	3,870	7,520	3,670	3,900	7,570
10-14	3,800	3,960	7,760	3,770	3,990	7,760	3,760	4,020	7,780	3,770	4,030	7,810
15-19	3,660	4,000	7,660	3,670	4,020	7,690	3,780	4,030	7,810	3,800	4,030	7,830
20-24	3,210	3,300	6,510	3,330	3,420	6,740	3,420	3,600	7,020	3,510	3,700	7,210
25-29	2,500	2,700	5,200	2,660	2,830	5,500	2,800	2,870	5,670	2,910	3,050	5,960
30-34	2,500	2,370	4,880	2,490	2,370	4,870	2,390	2,400	4,790	2,400	2,410	4,810
35-39	2,520	2,320	4,840	2,560	2,390	4,950	2,640	2,440	5,080	2,640	2,470	5,110
40-44	2,170	2,000	4,170	2,210	2,050	4,270	2,300	2,090	4,390	2,380	2,120	4,510
45-49	1,960	1,830	3,790	2,020	1,850	3,880	2,060	1,870	3,930	2,060	1,910	3,980
50-54	1,960	1,810	3,780	1,890	1,780	3,670	1,860	1,780	3,640	1,900	1,760	3,660
55-59	1,850	1,720	3,570	1,890	1,750	3,640	1,910	1,800	3,710	1,920	1,780	3,690
60-64	1,850	1,570	3,420	1,840	1,550	3,390	1,790	1,540	3,330	1,790	1,580	3,370
65-69	1,530	1,320	2,840	1,580	1,400	2,990	1,710	1,440	3,140	1,740	1,450	3,190
70-74	1,130	930	2,070	1,190	970	2,160	1,220	990	2,210	1,250	1,030	2,280
75-79	700	620	1,320	750	650	1,390	780	680	1,470	860	730	1,590
80-84	340	290	630	370	300	670	400	330	730	430	330	760
85+	260	160	410	260	170	440	280	190	460	270	200	470
All Ages	39,100	38,500	77,600	39,700	39,100	78,800	40,400	39,800	80,100	41,000	40,400	81,300



Age	Female	Male	Total									
Groups		2030			2031			2032			2033	
00-04	3,690	3,890	7,580	3,720	3,930	7,660	3,760	3,970	7,740	3,810	4,020	7,820
05-09	3,690	3,930	7,610	3,710	3,960	7,680	3,750	4,000	7,750	3,790	4,040	7,830
10-14	3,810	4,010	7,820	3,800	4,060	7,860	3,830	4,070	7,900	3,820	4,080	7,900
15-19	3,770	4,080	7,850	3,810	4,020	7,840	3,780	4,050	7,830	3,760	4,070	7,830
20-24	3,580	3,780	7,360	3,620	3,910	7,540	3,630	3,930	7,560	3,730	3,940	7,660
25-29	3,030	3,120	6,150	3,090	3,150	6,230	3,190	3,260	6,450	3,280	3,440	6,720
30-34	2,440	2,510	4,960	2,570	2,670	5,240	2,730	2,800	5,530	2,860	2,830	5,690
35-39	2,610	2,420	5,030	2,590	2,390	4,980	2,570	2,380	4,950	2,460	2,410	4,860
40-44	2,500	2,240	4,740	2,560	2,320	4,880	2,600	2,390	4,990	2,680	2,440	5,110
45-49	2,090	1,920	4,010	2,150	1,970	4,110	2,180	2,020	4,200	2,270	2,050	4,320
50-54	1,900	1,800	3,700	1,900	1,770	3,670	1,960	1,790	3,750	2,000	1,800	3,790
55-59	1,910	1,740	3,650	1,920	1,740	3,650	1,840	1,700	3,540	1,800	1,700	3,500
60-64	1,780	1,600	3,380	1,780	1,630	3,420	1,830	1,650	3,480	1,850	1,710	3,560
65-69	1,780	1,470	3,250	1,760	1,460	3,220	1,750	1,440	3,190	1,700	1,430	3,140
70-74	1,330	1,070	2,400	1,370	1,140	2,510	1,420	1,230	2,650	1,540	1,260	2,810
75-79	890	750	1,640	960	760	1,720	1,000	800	1,800	1,020	810	1,830
80-84	460	390	850	510	430	950	550	450	1,010	590	490	1,070
85+	290	210	490	300	210	510	330	220	550	350	240	590
All Ages	41,500	40,900	82,500	42,100	41,500	83,700	42,700	42,100	84,900	43,300	42,700	86,100



Age	Female	Male	Total									
Groups		2034			2035			2036			2037	
00-04	3,850	4,070	7,930	3,900	4,120	8,030	3,960	4,170	8,140	4,010	4,240	8,240
05-09	3,820	4,080	7,900	3,860	4,120	7,980	3,900	4,160	8,060	3,940	4,200	8,140
10-14	3,840	4,100	7,950	3,870	4,130	8,000	3,890	4,170	8,060	3,930	4,210	8,140
15-19	3,770	4,080	7,860	3,810	4,060	7,870	3,800	4,110	7,910	3,830	4,120	7,950
20-24	3,750	3,930	7,670	3,720	3,970	7,690	3,760	3,920	7,680	3,730	3,940	7,670
25-29	3,380	3,540	6,920	3,450	3,630	7,080	3,500	3,760	7,260	3,510	3,770	7,280
30-34	2,970	3,010	5,980	3,090	3,070	6,160	3,150	3,100	6,250	3,260	3,220	6,480
35-39	2,460	2,410	4,880	2,510	2,520	5,030	2,630	2,670	5,300	2,790	2,800	5,590
40-44	2,670	2,460	5,130	2,640	2,420	5,060	2,620	2,380	5,000	2,600	2,380	4,980
45-49	2,350	2,090	4,440	2,460	2,210	4,680	2,530	2,290	4,820	2,570	2,360	4,930
50-54	2,000	1,840	3,840	2,030	1,840	3,880	2,080	1,900	3,980	2,120	1,950	4,070
55-59	1,840	1,690	3,520	1,840	1,730	3,560	1,840	1,700	3,540	1,910	1,710	3,620
60-64	1,860	1,680	3,540	1,850	1,660	3,500	1,860	1,650	3,510	1,790	1,620	3,410
65-69	1,700	1,470	3,170	1,700	1,490	3,180	1,700	1,520	3,220	1,750	1,540	3,290
70-74	1,580	1,280	2,870	1,620	1,310	2,940	1,610	1,300	2,920	1,610	1,280	2,890
75-79	1,050	830	1,880	1,110	860	1,970	1,150	930	2,070	1,200	1,010	2,210
80-84	660	540	1,200	690	560	1,240	750	570	1,320	780	590	1,370
85+	360	250	610	390	280	690	430	320	750	480	330	810
All Ages	43,900	43,400	87,300	44,500	44,000	88,500	45,200	44,600	89,800	45,800	45,300	91,100



Age	Female	Male	Total									
Groups		2038			2039		2040			2041		
00-04	4,060	4,290	8,350	4,120	4,350	8,460	4,180	4,400	8,580	4,230	4,460	8,690
05-09	3,980	4,250	8,230	4,030	4,300	8,340	4,090	4,360	8,440	4,140	4,410	8,550
10-14	3,970	4,250	8,220	4,010	4,290	8,300	4,050	4,330	8,380	4,090	4,380	8,460
15-19	3,830	4,130	7,960	3,850	4,160	8,010	3,870	4,190	8,060	3,900	4,230	8,130
20-24	3,710	3,970	7,680	3,720	3,980	7,700	3,760	3,960	7,720	3,750	4,010	7,760
25-29	3,600	3,780	7,390	3,620	3,780	7,400	3,590	3,830	7,420	3,640	3,770	7,410
30-34	3,360	3,400	6,760	3,460	3,510	6,970	3,530	3,600	7,130	3,580	3,740	7,310
35-39	2,930	2,830	5,760	3,030	3,010	6,050	3,150	3,080	6,230	3,220	3,110	6,330
40-44	2,490	2,400	4,890	2,500	2,410	4,900	2,540	2,510	5,050	2,660	2,670	5,330
45-49	2,650	2,410	5,060	2,650	2,440	5,080	2,620	2,390	5,000	2,600	2,350	4,950
50-54	2,210	1,990	4,200	2,290	2,030	4,320	2,410	2,150	4,560	2,480	2,230	4,700
55-59	1,940	1,720	3,670	1,950	1,770	3,720	1,980	1,770	3,750	2,030	1,830	3,860
60-64	1,750	1,620	3,370	1,790	1,610	3,390	1,790	1,650	3,440	1,790	1,620	3,420
65-69	1,770	1,590	3,370	1,790	1,580	3,360	1,780	1,550	3,330	1,790	1,550	3,340
70-74	1,560	1,280	2,830	1,560	1,310	2,870	1,550	1,330	2,880	1,560	1,360	2,920
75-79	1,320	1,050	2,370	1,370	1,070	2,440	1,410	1,110	2,520	1,400	1,100	2,500
80-84	790	590	1,380	800	600	1,400	860	620	1,470	880	680	1,570
85+	520	370	890	580	420	990	620	460	1,070	690	470	1,160
All Ages	46,400	45,900	92,400	47,100	46,600	93,700	47,800	47,300	95,000	48,400	48,000	96,400



Ama Crauna	Female	Male	Total	Female	Male	Total
Age Groups		2042			2043	
00-04	4,280	4,510	8,800	4,340	4,580	8,920
05-09	4,190	4,470	8,670	4,250	4,530	8,780
10-14	4,130	4,420	8,550	4,170	4,470	8,640
15-19	3,940	4,270	8,210	3,980	4,310	8,290
20-24	3,780	4,020	7,800	3,780	4,030	7,810
25-29	3,600	3,790	7,400	3,580	3,810	7,400
30-34	3,590	3,750	7,340	3,690	3,770	7,460
35-39	3,330	3,230	6,570	3,430	3,420	6,850
40-44	2,820	2,790	5,620	2,960	2,830	5,780
45-49	2,580	2,350	4,930	2,470	2,370	4,840
50-54	2,520	2,300	4,820	2,600	2,350	4,940
55-59	2,070	1,880	3,950	2,160	1,920	4,080
60-64	1,860	1,640	3,500	1,890	1,650	3,550
65-69	1,720	1,520	3,240	1,690	1,530	3,220
70-74	1,610	1,380	2,990	1,640	1,440	3,080
75-79	1,400	1,080	2,470	1,360	1,080	2,430
80-84	930	760	1,690	1,040	790	1,840
85+	740	480	1,220	750	500	1,250
All Ages	49,100	48,700	97,800	49,800	49,400	99,100



Appendix 2: Technical notes

1. Explanation of statistical terms used in this report

95% confidence interval

Technical definition

A 95% confidence interval represents a range from a lower to an upper value that is likely to include the true average figure for the entire population. It suggests that if a similar sample of the total population was taken 100 times, the true value would be found within this range 95 times. This confidence interval can vary in size: a larger number of survey responses or participants, typically results in a narrower range, indicating more precise estimates, while a smaller number of responses may result in a broader range, indicating less certainty about the exact figure.

Plain English definition

When a health study gives a number, like how many people feel healthy, it's often not just one number but a range. This range is what's called a 95% confidence interval. It's like a safety net that says, 'We think the real number is in here.' And if we did the study over and over, 95 times out of 100, we'd get a number in this range. The more people we include in our sample, the smaller and more accurate this net becomes. So, if we ask only a few people, the net is wide, and we're less sure. If we ask a lot of people, the net gets tighter, and we're more sure we've got the right number.

Example from the report

In a survey assessing health status among residents of Te Moana a Toi⁹ (see table below), 13.0% of the sampled Māori population considered their health to be 'Excellent'. However, this percentage is an estimate from a sample of people in Te Moana a Toi, not the entire population. The 95% confidence interval, shown in brackets as "(9.8, 16.2)", indicates that there is a 95% probability that the actual percentage of all Māori residents who would rate their health as 'Excellent' falls within this range. If this survey were to be conducted 100 times with different sample groups, it is expected that 95 of those surveys would yield a true percentage that falls between 9.8% and 16.2%.

Table 6 - Health status reported by Māori aged 15 years and over, Te Moana a Toi, 2018

Health Status		Te Moana a Toi		Aotearoa			
	%	(959	% CI)	%	(95% CI)		
Excellent	13.0	(9.8,	16.2)	15.1	(14.0,	16.2)	
Very Good	40.2	(35.6,	44.9)	36.9	(35.4,	38.3)	
Good	30.1	(25.3,	35.0)	30.3	(29.0,	31.7)	
Fair/poor	16.6	(12.9, 20.3)		17.7	(16.6,	18.8)	

Source: Te Kupenga 2018, Statistics New Zealand customised report.

⁹ The example tables in this technical appendix are all taken from the Te Moana a Toi IMPB profile, and are presented purely as an example to facilitate understanding across all IMPB data profiles.



Age standardisation

Technical definition

Age-standardisation is a statistical method used to compare rates of events across different populations by adjusting for age differences in the two groups. This method is particularly useful when comparing health outcomes between groups like Māori and non-Māori, where there are significant differences in age distribution; for example only 8% of Māori are aged 65 and over in Te Moana a Toi compared with 26% of non-Māori (see the table below).

Because of these age differences, comparing crude rates (actual observed rates) can be misleading. By applying the age-specific rates from the populations being compared to a standard population, age-standardised rates provide a clearer comparison as if the populations had the same age distribution. Almost all data in this report has been age-standardised to the 2001 Māori population. Where crude rates are presented instead, this is noted beneath the table.

Table 2 - Population estimate by age group, Te Moana a Toi, 2023

Age group (years)		Māori		non-N	Total IMPB	
	Number	Age distribution	% of IMPB	Number	Age distribution	number
0–14	20,255	30%		30,670	15%	50,925
15–24	12,285	18%		16,810	8%	29,095
25-44	16,465	24%		50,870	25%	67,335
45–64	13,030	19%		52,935	26%	65,965
65+	5,575	8%		51,760	26%	57,335
Total	68,000	100%	25%	202,740	100%	270,740

Plain English definition

Age-standardisation is a method used to compare health between two groups fairly. It adjusts the numbers to consider how young or old the people in each group are. This way, when looking at health data, it is more likely that any differences between the groups are not just because one has more young people or more old people. It helps give a more accurate picture of health when comparing two groups with a different spread of ages.

Example from the report

The table below shows an age-standardised rate of 28.4 per 100,000 per year ischaemic heart disease events among Bay of Plenty DHB Māori women between 2014 and 2018. Without age standardisation calculations, crude rates would be lower than 28.4 among Māori women. The lower rate would be simply because a larger proportion of the Māori population is younger and ischaemic heart disease is more frequent in older people.

Table 6 - Leading causes of death for Māori, all ages, Bay of Plenty DHB, 2014 to 2018

	Māori			non-Māori					
Cause	Av. no. per year	rate	-standardised per 100,000 (95% CI)	Av. no. per year		-standardised e per 100,000 (95% CI)	Māori/non-Māori rate ratio (95% CI)		Rate difference
Female									
Ischaemic heart disease	19	28.4	(16.2, 45.5)	98	8.3	(6.2, 10.9)	3.40	(1.95, 5.93)	20.1



Rate ratios

Technical definition

Rate ratios, often referred to as relative risks, are a measure of the relationship between the occurrence of a certain event in two different groups, typically standardised for age (see section on age standardisation above) to allow fair comparison. It is the result of the rate of the event in the first group (for example, Māori) divided by the rate in the second group (non-Māori), which serves as the reference group. A rate ratio of 1 indicates parity between groups, above 1 indicates a higher rate in the first group, and below 1 indicates a lower rate. In general, the data presented in this report uses Māori as the first group and compares it with non-Māori as the second group.

Plain English definition

A rate ratio compares how common something, like a disease, is between two different groups of people, like Māori and non-Māori. If the ratio is exactly 1, both groups are equally affected. If it's higher than 1, it means that the first group, in this case Māori, has the event happen more often. If it's lower, Māori have it happen less often. It tells us the relative disparity between two groups.

Example from the report

In the table below, the rate ratio for ischaemic heart disease is 3.40. This tells us that Māori females are more than three times as likely to suffer from this condition compared to non-Māori females after considering the age distribution in each group.

The 95% confidence interval (see section on confidence intervals above) of 1.95 to 5.93 for this rate ratio indicates that we are very sure that the true rate ratio is significantly different from 1, indicating a genuine disparity in risk between the two populations. In this report, a statistically significant difference between groups is evident when the confidence interval for the rate ratio does not cross 1. These results are shown in **bold** type.

Table 6 - Leading causes of death for Māori, all ages, Bay of Plenty DHB, 2014 to 2018

	Māori			non-Māori						
Cause	Av. no. per year	Age-standardised rate per 100,000 (95% CI)		Av. no. per year	Age-standardised rate per 100,000 (95% CI)		Māori/non-Māori rate ratio (95% CI)		Rate difference	
Female										
Ischaemic heart disease	19	28.4	(16.2, 45.5)	98	8.3	(6.2, 10.9)	3.40	(1.95, 5.93)	20.1	



Rate difference

Technical definition

Rate differences, also known as absolute differences, quantify the disparity between two groups by showing the additional number of events occurring in one group compared to another, per population unit (like per 100,000 people). This is calculated by subtracting the event rate of the reference group from that of the comparison group.

Plain English definition

Rate difference tells us how much more often something happens in one group compared to another. If you take the number of times an event happens per 100,000 people in one group and subtract the same from another group, you get the rate difference. This number shows if one group is experiencing more of a certain event, like a disease or death, and by how much. It's a simple way to see the actual impact of a problem on one group over another.

Example from the report

The table below show that Māori females in Bay of Plenty DHB have an age-standardised rate of ischaemic heart disease at 28.4 events per 100,000 per year, while the rate for non-Māori females is 8.3. This gives a rate difference of 20.1 events per 100,000 per year, which tells us that in a population of 100,000 Māori women and 100,000 non-Māori women there are 20.1 more cases of ischaemic heart disease among Māori females than non-Māori females each year. This figure is crucial because it doesn't just show the relative disparity (like a rate ratio does), but it tells us how many additional events are affecting Māori females, highlighting the actual impact of the disease on the population and where health resources might be most needed to address the disparity.

Table 6 - Leading causes of death for Māori, all ages, Bay of Plenty DHB, 2014 to 2018

Cause	Māori			non-Māori					
	Av. no. per year	rate	-standardised per 100,000 (95% CI)	Av. no. per year	-	e-standardised e per 100,000 (95% CI)	Māori/non-Māori rate ratio (95% CI)		Rate difference
Female									
Ischaemic heart disease	19	28.4	(16.2, 45.5)	98	8.3	(6.2, 10.9)	3.40	(1.95, 5.93)	20.1



2. Key methods and quality limitations of key data sources

This section describes in more detail the specific methods, and key limitations, used for each of the main data sources used in this report.

Numerators

Data in this first volume of IMPB profiles are sourced from Te Whatu Ora, Manatū Hauora (the Ministry of Health), and Statistics New Zealand (StatsNZ). Where administrative data (e.g. national mortality data) are used, the most recent five years of non-provisional data were aggregated to provide more stable rate estimates for smaller areas. Census data were taken from the 2018 Census, and data from the Te Kupenga survey were from the 2018 Te Kupenga survey, undertaken after the 2018 Census.

Denominators

StatsNZ mid-year (at 30 June) estimated resident population was used as denominator data in the calculation of population rates for deaths and Primary Healthcare Organisation (PHO) enrolment. For census variables, the denominator is the people for whom there is a response / relevant information from the census dataset for the question asked ('people stated'). This differs for each question, and is a subset of the total usually resident population identified by the census for the relevant rohe (region). For Te Kupenga survey data, the denominator is the total stated population, this means that people who refuse to answer/ don't know their answer/ answer with an invalid answer are excluded.

Ethnicity data

Ethnicity data quality

Although high quality ethnicity data are critical for Māori health improvement, ethnicity data quality in the health sector remains poor (Harris, Paine et al. 2022). It is the responsibility of the entire health system to collect, record and report ethnicity data in the ways set out in the HISO 10001:2017 Ethnicity Data Protocols (Ministry of Health. 2017). Despite the protocols being in existence for nearly 20 years, there is evidence that they are not being adhered to and Māori have continued to be systematically undercounted (Cormack D and McLeod M 2010, Harris, Paine et al. 2022). Self-identified ethnicity recorded on the Census is considered to be the "gold-standard" for ethnicity data, so this is used as the denominator for most variables in this report.

To understand what impact the ethnicity data quality is likely to have, on the accuracy of the results presented in this report, we need to consider the ethnicity data quality in both the numerator and the denominator. For some measures, it may underestimate the true number of, or rate of, a particular outcome for Māori. The potential impact of ethnicity data weaknesses is discussed for each data source later in this Appendix.

Ethnicity classification

When analysing data, there are different ways to classify people who report multiple ethnicities. The two main ways are *total response* (*overlapping*) *output* and *prioritised output*. In total response output, each respondent is counted in each of the ethnic groups they reported. So, individuals who indicate more than one ethnic group are counted more than once, and the sum of the ethnic group populations will exceed the total population of NZ. For example, using total response classification, a death from lung cancer in an individual who identifies as Māori and New Zealand European, will be reported as a lung cancer death for both ethnicities.

In prioritised output, each respondent is allocated to a single ethnic group using a prioritisation order, with Māori first, to ensure that ethnic groups of policy importance or of small size, are not swamped by the New Zealand European ethnic group. Under this method, a person is classified as Māori if any

one of their recorded ethnicities are Māori. For example, using prioritised classification, a death from lung cancer in a person recorded as both Māori and New Zealand European, would be counted as a lung cancer death for Māori, and not in non-Māori.

In this report, the method of ethnicity classification is noted under each table or figure. Wherever possible, prioritised ethnicity classification was used when people identified with more than one ethnic group.

Comparison group

Most indicators compare Māori with non-Māori. Non-Māori includes all people who do not identify as Māori and represent a comparative or reference group. Some indicators in this report (e.g. life expectancy) use non-Māori non-Pacific (all people who do not identify as either Māori or Pacific or both) as the comparison group. This is done because in areas where there are large Pacific populations, grouping the Pacific population with the non-Māori group skews the result for the comparison group toward the Māori population. This is particularly necessary in regions where there is a high Pacific population such as South Auckland.

Age-standardised and crude rates

This report uses direct age-standardisation; most rates (unless noted otherwise) are standardised to the 2001 Census Māori population. Where data were not available with sufficient age group breakdown to allow age standardisation, or data for a specific age were presented, crude rates were calculated. In this case, caution should be taken when comparing Māori with non-Māori results. Crude rates accurately portray a situation in each population, but make comparisons difficult, because they do not consider the different age distributions in each of the populations (e.g., the Māori population is much younger than the non-Māori population). Rates were not calculated for counts fewer than five in data from national collections. For Te Kupenga data, if the weighted count (estimate) was less than 1000 then the data was supressed.

Confidence intervals

This report has endeavoured where possible to provide local data specific to IMPBs and their relevant DHB areas. Some of these areas have small populations. As the size of the group becomes smaller, the confidence interval (CI) becomes wider, and there is less certainty about the rate. This means the degree of confidence and certainty about the numbers diminishes for rohe (regions) with smaller populations. Thinking of the data as 'indicative' rather than precise is important in these rohe, as well as considering Māori-specific regional and national data, which will have greater certainty around rates, because of the larger sample size.

When the CIs of two groups do not overlap, the difference in rates between the groups is considered statistically significant. Sometimes, even when there are overlapping CIs, the difference between the groups may be statistically significant. Determining that would require further statistical testing which has not been undertaken for this report.

Rate ratios

Age-standardised rate ratios are used in this report to compare age-standardised rates between Māori and non-Māori. The rate ratio (RR) is equal to the age-standardised Māori rate divided by the age-standardised non-Māori rate. The non-Māori population is used as the reference population. For example, an age-standardised RR of 1.5 means that the rate is 50 percent higher (or 1.5 times as high) in Māori than in non-Māori, after taking into account the different age structures of these two populations. This report gives rate ratios and their 95 percent Cls. In this profile, if the Cl of the rate ratio does not include the number 1, the ratio is said to be statistically significant. Differences presented in this profile in **bold** are statistically significant.

Demography data

Indicators on population demography and projections use the estimated resident population (ERP) and projections provided by StatsNZ for the health sector, from a 2018 base. The ERP is an estimate designed to adjust for the undercount for various groups in the census response rate, people temporarily overseas or elsewhere in NZ from their usual residence on census night, and key population changes (births, deaths, mobility) since the 2018 census.

In the estimates and projections prioritised ethnicity was used to identify Māori individuals (any person who identified Māori as any of their ethnic groups in the base census data on which the estimates and projections are built) and non-Māori included people who had at least one valid ethnic response, none of which was/were Māori.

The Census of Population and Dwellings

Indicators using data from the 2018 Census of Population and Dwellings are derived from the census usually resident (UR) population (residents of an area living in the area on census night and people living elsewhere in Aotearoa from their usual residence on census night). Data used in this report were sourced from the publicly available UR data provided on the StatsNZ website, and for some indicators, from a custom data extract produced by StatsNZ for the previous Northern Region DHBs (which included data for the whole of Aotearoa).

StatsNZ apply confidentiality rules to census data to protect the confidentiality of individuals, families, households, dwellings, and undertakings in 2018 Census data. Counts are calculated using a method called fixed random rounding to base 3, and suppression of 'sensitive' counts less than six, where tables report multiple geographic variables and/or small populations. This means individual figures may not always sum to stated totals¹⁰.

Due to changes in the 2018 Census methodology and lower than anticipated response rates, as described further below, time series data for census variables should be interpreted with care.

Most census variables in the Wai Ora chapter have been age-standardised to the 2001 Māori population. The unpaid work variables were not able to be age-standardised for this report, and crude rates are presented. In this case, caution should be taken when comparing Māori with non-Māori results.

The 2018 Census was the first 'digital-first' census undertaken in Aotearoa, as a part of modernising and streamlining the census process. Unfortunately, the 2018 Census had a very low response rate overall, and especially for Māori and Pacific peoples - approximately 68% for Māori and 65% for Pacific peoples. Adjustments were made to improve the quality of the data (for example, using data from previous censuses and other administrative datasets), and the overall quality of the 2018 Census data is now considered moderate/good. However, the adjustments do not affect the Māori and non-Māori population in the same way. For example, in the 2018 Census, 29% or more of the ethnicity data for Māori came from other sources. This means that the ethnicity data in the 2018 census for Māori is not of the same quality as the data for the NZ European ethnic population, for example, which had only 11.5% of their responses from these other sources.

Further details on the adjustment methods used in the 2018 Census can be found online via Stats NZ¹¹. In summary, the core self-response data from the 2013 Census was combined with administrative data (e.g. from the education or health system), and in some situations data derived by statistical models to predict what the missing data would have been (called imputation). In addition

https://www.stats.govt.nz/assets/Uploads/Reports/Final-report-of-the-2018-Census-External-Data-Quality-Panel/Downloads/Final-report-of-the-2018-Census-External-Data-Quality-Panel-corrected.pdf



¹⁰ More info on Census confidentiality rules: <u>Applying confidentiality rules to 2018 Census data and summary of changes since 2013</u> |
Stats N7

to different levels of self-response, people identified as living in NZ at the time of the census have different levels of information from other sources available to StatsNZ to draw on.

However, on the other hand, the census is a key source for population level data about factors that are important for health, such as income, employment, and housing. StatsNZ has provided quality ratings for the 2018 Census data to help users determine how to interpret the data. Along with StatsNZ's own quality ratings, they also engaged an External Data Quality Panel which included Māori population experts, who provided their assessment of the census data quality. The table below shows the ratings of both for the data variables used in this report. The overall message from these ratings is that the data can provide insights into the situation for Māori whānau, but it should be seen as indicative, rather than precise.

Table 34 - Quality ratings 2018 Census variables included in this report

Variable name	StatsNZ quality rating	External Data Quality Panel quality rating	Notes
Census usually resident population count	Very high	Very high	
Ethnicity	High	Moderate	
Number of bedrooms	High	High	Number of bedrooms is used to help derive estimates of household crowding. There were over 300,000 people who could not be placed into households in the 2018 data. This means the number of people who lived in a crowded house may be undercounted.
Number of rooms	Moderate	Poor	
Housing quality: dwelling dampness and mould indicators	Moderate	Moderate	This is a self-evaluated assessment of whether the home has mould that is larger than an A4 sheet of paper (in total).
Main types of heating and fuel types used to heat dwellings	Moderate	Moderate	This question was first introduced in the 2018 Census. Each type of heating reported was recorded once only.
Tenure of household	Moderate	Moderate	
Access to telecommunication systems	Moderate	Moderate	The online data collection methodology of the 2018 Census may have affected this variable. The proportion of households with no access to telecommunications was lower than expected. The proportion of households with access to a telephone was higher than expected. This data provides information on access to telecommunication systems at the household level. It does not show whether a particular household member has access to those amenities. In some cases, not every member of a household has equal access to particular telecommunication systems.
Number of motor vehicles	Moderate	Moderate	
<u>Industry</u>	High	High	Industry is the type of activity undertaken by the organisation or business where people work.
Occupation	Moderate	Poor	An occupation is a set of jobs that require the performance of similar or identical sets of tasks. Occupations are organised based on skills, using the ANZSCO classification. The significant use of imputation may have

Variable name	StatsNZ quality rating	External Data Quality Panel quality rating	Notes
			inflated the total number of respondents in all categories.
Qualifications: highest qualification	Moderate	Moderate/poor	
Total personal income	High	High	Total personal income received is the total before-tax income of a person in the 12 months ended 31 March 2018. The information is collected as income bands rather than in actual dollars. This includes all possible sources of income.
Status in employment	High	Moderate	Employment is described as full-time (30 hours or more / week) or part-time (< 30 hours per week). A person not employed is described as either 'unemployed' or 'not in the labour force'. Not in the labour force means not employed and not actively seeking work or not available for work
Unpaid activities	Poor	Not applicable	Because of the low quality ratings, Stats NZ recommend very careful use of this data particularly for Māori and Pacific peoples and at small geographies. No alternative data source or imputation was available to replace missing responses.

Geographical alignment between IMPB and DHB areas

This report has endeavored to report data specific to each IMPB health planning area and has used several slightly different methods to do this in different chapters of the report.

For population estimates, and Te Kupenga survey data, the population for an IMPB has been calculated using geographies (SA2 areas or Territorial Authority/Local Boards) that are smaller than the previous DHB districts, to be able to better align with the IMPB health planning areas. This means the Te Taura Ora o Waiariki and Tūwharetoa IMPBs have been able to be split out separately, and Ōtāhuhu has been included as part of Ngaa Pou Hauora oo Taamaki Makaurau, rather than Te Taumata Hauora o Te Kahu o Taonui (historically Ōtāhuhu was part of Auckland DHB rather than Counties Manukau DHB, so the Auckland Council Local Board Māngere-Ōtāhuhu spanned the boundary between the DHBs)¹². In some cases, for example at the Nelson-Marlborough/Te Tauraki border, the IMPB health planning area did not align completely with SA2 areas.

There may be some variation between the IMPB population estimates presented here compared to estimation using data from the previous DHB. This is due to there being a higher level of uncertainty around the SA2 population estimates and they will not always sum to exactly the same population by age, sex and ethnicity as the district population estimates.

For other measures, including mortality data, NZDep2018 and PHO enrolment, the IMPB population has been calculated using the sum of the main DHBs it contains. So, for example IMPB mortality data for Te Taumata Hauora o Te Kahu o Taonui will include all of Northland, Auckland and Waitematā DHBs, even though that includes communities such as Ōtāhuhu which are not part of the IMPB.

¹² Ōtāhuhu has a population of approximately 16,000 people, the majority of whom identify as Pacific and Asian (Indian). The area is classified as NZDep2018 deciles 9 and 10 – the most socio-economically challenged areas.

Life expectancy

There are two parts to the life expectancy data provided in this report. There is a 'standard' calculation of life expectancy at birth for each IMPB, using mortality data from Manatū Hauora and population data from StatsNZ and presented as the gap between Māori and non-Māori. It uses five years of data to be able to provide ethnicity and male/female information.

There is also information on what conditions contribute to those life expectancy gaps, from an analysis completed by the Service Innovation and Improvement Directorate, Te Whatu Ora in May 2023 titled "The Contribution of Avoidable Mortality to the Life Expectancy Gap among the Māori and Pacific population. Regional Summary." This analysis compared Māori with the non-Māori, non-Pacific population, so that is why this comparator group is used for this section in this IMPB report.

The Arriaga method—a life table decomposition technique accounting for both age and cause of death—was used. The analyses and calculations are based on official death data from the Te Whatu Ora mortality collection, while population data are derived from official StatsNZ population estimates.

The analysis hinges on the principal underlying cause of death classification, which simplifies the reality that multiple factors can contribute to a single death. This may result in an underestimation of the effects of prevalent conditions contributing to, but not the final causes of death. As it requires cause of death information, these are often two years delayed to allow coronial processes to be completed. As such, the life expectancy figures here may not be the most recent available, but are the most recent that allows this type of gap analysis.

Causes of death are divided into 50 potentially avoidable conditions. Avoidable deaths encompass those deemed amenable to high-quality healthcare, preventable through public health interventions, or both. A comprehensive list of the conditions used in this analysis, along with their corresponding ICD codes, can be found in the Te Whatu Ora report. Most are limited to those under 75 years, except leukemia which is only considered avoidable under the age of 45 years and external injuries which includes all ages.

Mortality data

Indicators on cause of death and mortality come from the national Mortality Collection. This classifies the underlying cause of death for all deaths registered in Aotearoa and all registered fetal deaths (stillbirths). Aotearoa is currently using the International Statistical Classification of Diseases and Related Health Problems, Tenth Revision, Australian Modification (ICD-10-AM) classification and the World Health Organization (WHO) ICD Rules and Guidelines for Mortality Coding. Mortality data are presented for Māori and non-Māori. In each data set a person was classified as Māori if any one of their recorded ethnicity was Māori. The year range of 2014 to 2018 was used as complete mortality data records were not available for 2019 and 2020 at the time of writing. The DHB of residence was determined from the domicile code attached to the death registration (so even if a person passed away at a tertiary hospital outside their home region, their death would be recorded as one in their home DHB). In tables presenting data on causes of death, data is not presented where there were fewer than five Māori events during the period represented by the data. There are several different methods of classifying causes of death as "potentially avoidable", "preventable" or "amenable". The ICD-10-AM codes used for potentially avoidable death tables in this report are listed in the next Appendix.

Te Kupenga Survey

Te Kupenga 2018 is StatsNZ's survey of Māori wellbeing. A survey of almost 8,500 adults (aged 15 years and over) of Māori ethnicity and/or descent, Te Kupenga gives an overall picture of the social, cultural, and economic wellbeing of Māori people in Aotearoa.

Te Kupenga is a post-census survey. This means the survey sample was selected from people who identified as having Māori ethnicity and/or descent on their 2018 census form, so only those who

completed the census were able to be selected. Given that a lower proportion of Māori people completed the 2018 Census than planned or anticipated, StatsNZ investigated the potential impact this may have had on the Te Kupenga sample. They found some bias in the sample frame (the group of people who could have been selected to participate) compared with the total Māori population. However, this bias was small, and they were able to remove most of the effect of the bias through the statistical weighting process. See StatsNZ website for more information on this¹³.

In this IMPB profile, all estimates of numbers, percentages, and confidence intervals for data presented from Te Kupenga were calculated by StatsNZ and provided in a customised extract. Estimates of counts were rounded to the nearest thousand. Estimates of proportions were rounded to 1 decimal point. All percentages were calculated from unrounded data. If the weighted count (estimate) was less than 1000 then the data was supressed. Further details on the survey measures are available in the Te Kupenga 2018 report and can be found at the StatsNZ website 14.

Primary care enrolment

Primary care enrolment data is based on the National Enrolment System using the National Health Index (NHI). Ethnicity data in the NHI is known to undercount Māori by 15.7% compared to the ethnicity people report in the census, with higher undercounts for Māori men (Harris, Paine et al. 2022). The denominator for calculating the percentage of people enrolled in a PHO is the estimated resident population, which uses ethnicity based on the 2018 Census. The poor ethnicity data quality in the NHI makes it difficult to assess how many Māori are actually missing out on being enrolled with primary health care, and how many are actually enrolled but misclassified with a non-Māori ethnicity. It is likely that both of these factors make a contribution to the inequity in primary care enrolment data. Primary care enrolment data presented in this report are not age-standardised. In this case, caution should be taken when comparing Māori with non-Māori results. Crude rates make comparisons difficult, because they do not take into account different age distributions in each of the populations.

NZ Index of Deprivation 2018

NZDep2018 is an area-based measure of relative socio-economic deprivation. It is based on nine variables from the 2018 Census which cover eight different dimensions of socio-economic hardship. These variables relate to home internet access, receipt of welfare benefits, household income, employment, qualifications, home ownership, family structure, household crowding and housing quality. NZDep2018 gives a deprivation score for small area geographies (i.e. meshblocks, and SA1s) (Atkinson, Salmond et al. 2019). These scores are aggregated into deciles (1-10, 1 being areas with the least socio-economic challenge and 10 being those the most disadvantage). This report uses NZDep2018 information supplied by StatsNZ for the health sector, applying the scores to estimated resident populations to estimate the number of people living in each decile.

Geographic Classification of Health

The Geographic Classification for Health (GCH) is a rural-urban geographic classification designed to allow Aotearoa's health researchers and policy makers to accurately monitor rural-urban variations in health outcomes. The GCH classifies all areas of Aotearoa as rural or urban according to their proximity to larger urban areas with respect to health (Whitehead, Davie et al. 2021).

The GCH is composed of five categories, two urban and three rural, that reflect degrees of reducing urban influence and increasing rurality. 'Urban 1' to 'Urban 2' are based on population size, and 'Rural 1' to 'Rural 3' based on drive time to their closest major, large, medium, and small urban areas. The

¹⁴ https://www.stats.govt.nz/information-releases/te-kupenga-2018-final-english/



¹³ https://www.stats.govt.nz/methods/assessment-of-potential-bias-in-the-te-kupenga-sample-frame-2018

population and drive time thresholds used in the GCH were developed from a health perspective and tested in partnership with a wide range of rural health stakeholders.



Appendix 3: ICD-10-AM Codes

The International Classification of Diseases (ICD-10-AM) codes used for the calculation of potentially avoidable mortality are presented below.

Table 35 - Potentially avoidable mortality ICD-10-AM codes

Condition	ICD-10-AM Code
Tuberculosis	A15-A19, B90
Selected invasive bacterial and protozoal infection	A38-A41, A46, A481, B50-B54, G00, G03, J020, J13-J15, J18, L03
Hepatitis	B15-B19
HIV/AIDS	B20-B24
Lip, oral cavity and pharynx cancers	C00-C14
Oesophageal cancer	C15
Stomach cancer	C16
Colorectal cancer	C18-C21
Liver cancer	C22
Lung cancer	C33-C34
Melanoma of skin	C43
Non-melanotic skin cancer	C44
Breast cancer (female only)	C50
Cervical cancer	C53
Uterine cancer	C54-C55
Bladder cancer	C67
Thyroid cancer	C73
Hodgkin's disease	C81
Leukaemia	C910-C911
Benign tumours	D10-D36
Thyroid disorders	E00-E07
Diabetes	E10-E14
Alcohol-related diseases	F10, I426, K292, K70
Illicit drug use disorders	F11-F16, F18-F19
Epilepsy	G40-G41
Birth defects	H311, P00, P04, Q00-Q99
Rheumatic and other valvular heart disease	101-109
Hypertensive heart disease	110-115
Nephritis and nephrosis	I12-I13, N00-N09, N17-N19

TE MOANA A TOI HEALTH PROFILE VOLUME ONE

Condition	ICD-10-AM Code
Ischaemic heart disease	120-125
Deep vein thrombosis with pulmonary embolism	126, 1802
Cerebrovascular diseases	160-169
Aortic aneurysm	171
Viral pneumonia and influenza	J10, J12, J171, J21
COPD	J40-J44
Asthma	J45-J46
Peptic ulcer disease	K25-K28
Acute abdomen, appendicitis, intestinal obstruction, cholecystitis/lithiasis, pancreatitis, hernia	K35-K38, K40-K46, K80-K83, K85-K86, K915
Chronic liver disease (excluding alcohol-related disease)	K73-K74
Obstructive uropathy and prostatic hyperplasia	N13, N20-N21, N35, N40, N991
Complications of perinatal period	P03, P05-P95
Motor vehicle accidents	V01-V04, V06, V09-V80, V87, V89, V99
Falls	W00-W19
Drownings	W65-W74
Fires, burns	X00-X09
Accidental poisonings	X40-X49
Suicide and self-inflicted injuries	X60-X84, Y870
Violence	X85-Y09, Y871



Appendix 4: Māori 2001 Population

The table below shows the 2001 Māori population standard used for age-standardisation in this report, including the weightings applied to each age-group.

Table 36 - 2001 Census total Māori population

Age group (years)	2001 Census total Māori population	Weighting
0-4	67,404	12.81
5-9	66,186	12.58
10-14	62,838	11.94
15-19	49,587	9.42
20-24	42,153	8.01
25-29	40,218	7.64
30-34	39,231	7.46
35-39	38,412	7.30
40-44	32,832	6.24
45-49	25,101	4.77
50-54	19,335	3.67
55-59	13,740	2.61
60-64	11,424	2.17
65-69	8043	1.53
70-74	5046	0.96
75-79	2736	0.52
80-84	1251	0.24
85+	699	0.13



Te rārangi tohutoro References

Te rārangi tohutoro - References

Atkinson, J., C. Salmond and P. Crampton (2019). "NZDep2018 Index of Deprivation, interim research report." <u>University of Otago: Wellington, New Zealand</u> **5541**: 1-65.

Cormack D and McLeod M (2010). <u>Improving and maintaining quality in ethnicity data collections in</u> the health and disability sector. Wellington, Te Rōpū Rangahau Hauora a Eru Pōmare.

Curtis, E. (2016). "Indigenous positioning in health research: the importance of Kaupapa Māori theory-informed practice." <u>AlterNative</u>: An International Journal of Indigenous Peoples **12**(4): 396-410.

Curtis E, Loring B, Harris R, McLeod M, Mills C, Scott N and Reid P (2022). Māori Health Priorities. A report commissioned by the interim Māori Health Authority (iMHA) to inform development of the interim New Zealand Health Plan (iNZHP). Auckland, Te Aka Whai Ora.

Curtis, E., R. Jones, E. Willing, A. Anderson, S.-J. Paine, S. Herbert, B. Loring, G. Dalgic and P. Reid (2023). "Indigenous adaptation of a model for understanding the determinants of ethnic health inequities." <u>Discover Social Science and Health</u> **3**(1): 10.

Harris, R., S. J. Paine, J. Atkinson, B. Robson, P. T. King, J. Randle, A. Mizdrak and M. McLeod (2022). "We still don't count: the under-counting and under-representation of Māori in health and disability sector data." N Z Med J **135**(1567): 54-78.

Ministry of Health. (2017). HISO 10001:2017 Ethnicity Data Protocols. Wellington, Ministry of Health.

Simmonds, S., B. Robson, F. Cram and G. Purdie (2008). "Kaupapa Māori Epidemiology." Australasian Epidemiologist., **15**(1): 2-6.

Whitehead, J., G. Davie, B. de Graaf, S. Crengle, M. Smith, R. Lawrenson, D. Fearnley, N. Farrell and G. Nixon (2021). "The Geographic Classification for Health." <u>Methodology and classification</u> report.



